

# REXOM<sup>TM</sup> BROMHEXINE ELIXIR

Bromhexine Hydrochloride (4mg/5ml)

## What is in this leaflet

1. What REXOM<sup>TM</sup> Bromhexine is used for
2. How REXOM<sup>TM</sup> Bromhexine works
3. Before you use REXOM<sup>TM</sup> Bromhexine
4. How to use REXOM<sup>TM</sup> Bromhexine
5. While you are using REXOM<sup>TM</sup> Bromhexine
6. Side effects
7. Storage and disposal of REXOM<sup>TM</sup> Bromhexine
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of revision

## What REXOM<sup>TM</sup> Bromhexine is used for

It is used to loosen the mucus, making it easier to cough up the mucus for airway diseases, accompanied by excessive mucus and impaired mucus clearance.

## How REXOM<sup>TM</sup> Bromhexine works

REXOM<sup>TM</sup> Bromhexine helps to clear the chest by thinning the mucus in your air passages.

## Before you use REXOM<sup>TM</sup> Bromhexine

### -When you must not use it

It should not be used if you had known to be hypersensitive to Bromhexine or other components of the formulation.

### -Before you start to use it

- There will be an expected increase in the flow of secretions.
- Should be used with care in people with a history of ulcer in digestive tract and asthma.

### *Pregnancy and lactation*

Talk to your doctor before taking this medicine if you are pregnant, might become pregnant or are breastfeeding. Bromhexine should only be used if the maternal condition justifies the potential risk to the fetus or baby.

### -Taking other medicines

Inform your doctor or pharmacist before taking this medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and REXOM<sup>TM</sup> Bromhexine may interfere with each other. They may be affected by REXOM<sup>TM</sup> Bromhexine or may affect how well it works.

- It may increase the concentration of concurrently used antibiotics.

## How to use REXOM<sup>TM</sup> Bromhexine

### -How much to use

Always use REXOM<sup>TM</sup> Bromhexine exactly as your doctor has prescribed. You should check with your doctor or pharmacist if in doubt.

### The usual dose:

Under 2 years: 1.25 ml  
2 to 6 years: 2.5 ml  
6 to 12 years: 5 ml

### -When to use it

To be taken 3 times a day. Use as directed by your doctor or pharmacist.

### -How long to use it

Continue taking REXOM<sup>TM</sup> Bromhexine as your doctor recommended. If symptoms persist, consult your doctor.

### -If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

### -If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

## While you are using REXOM<sup>TM</sup> Bromhexine

### -Things you must do

- Tell all the doctors, dentists and pharmacists treating you that you are taking REXOM<sup>TM</sup> Bromhexine.
- Tell your doctor immediately if you become pregnant while taking this medication.

### -Things you must not do

- Do not take more than the recommended dose unless your pharmacist or doctor tells you.
- Do not take any new medicines without consulting your doctor or pharmacist.
- Do not give REXOM<sup>TM</sup> Bromhexine to anyone else, even if they have the same symptoms or condition as you.

### -Things to be careful of

Be careful driving or operating machinery until you know how REXOM<sup>TM</sup> Bromhexine affects you.

## Side effects

Like all medicines, REXOM<sup>TM</sup> Bromhexine can cause side effects, although not everybody gets them.

If side effects persist, do not take the medicine and refer to the doctor.

REXOM<sup>TM</sup> Bromhexine may cause severe allergy and serious skin reactions. Stop using REXOM<sup>TM</sup> Bromhexine and seek medical assistance immediately if you experience any of the following symptoms:

- 1) severe allergy: breathing difficulties, light headedness, skin swellings or rash.
- 2) severe skin reaction: skin reddening, blisters, rash, fever, sore throat or eye irritation.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website [npra.moh.gov.my](http://npra.moh.gov.my) [Public → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

## Storage and disposal of REXOM<sup>TM</sup> Bromhexine

### -Storage

Keep container tightly closed. Store below 30°C. Protect from heat and light. Keep medicine out of reach of children.

### -Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

# REXOM™ BROMHEXINE ELIXIR

Bromhexine Hydrochloride (4mg/5ml)

---

## **Product description**

### -What it looks like

Colourless liquid with strawberry flavor.

### -Ingredients

-Active ingredient

Each 5ml contains

Bromhexine Hydrochloride (4mg)

-Inactive ingredients

Strawberry flavour, Menthol Crystals,  
Sodium Benzoate, Sodium Citrate,  
Saccharin Sodium, Citric Acid  
Anhydrous, Propylene Glycol,  
Hydroxypropyl Methylcellulose and  
Purified Water.

### -MAL number

MAL19962479XZ

## **Manufacturer and Product Registration Holder**

KCK Pharmaceutical Industries Sdn.  
Bhd.

Plot 61, Bayan Lepas Industrial Park  
Phase 4, Lintang Bayan Lepas 1,  
11900 Bayan Lepas, Pulau Pinang,  
Malaysia.

## **Date of revision**

16/06/2021

## **Serial Number**

NPRA (R2/1) 210521/682