


15mm

48mm

112mm

30mm



REXOM™
UNIZET
SYRUP

REXOM™
UNIZET
SYRUP 100 ml

REXOM™
UNIZET
SYRUP

REXOM™
UNIZET
SYRUP 100 ml

Keep the bottle tightly closed
protected from light.
Store below 30°C.
"Controlled Medicine"/ "Ubat Terkawal"
MAL20026062AZ



Each 5 ml contains :
Diphenhydramine Hydrochloride 8 mg
Bromhexine Hydrochloride 4 mg
Ammonium Chloride 100 mg
Sodium Benzoate 5 mg
(as preservative)

Dosage :
Adults : 10 ml (2 teaspoonfuls) of syrup
3 times a day, not to exceed 6 tsp in 24 hours.
Children aged 6-12 years : 5 ml (1 teaspoonful)
of syrup 3 times a day. Maximum daily dose :
15 ml syrup.

Not to be used in children less than 2 years of age.
To be used with caution and doctor's/
pharmacist's advice in children 2 to 6 years of
age.

Warning :

When used for the treatment of cough
and cold:
(a) Not to be used in children less than 2
years of age.
(b) To be used with caution and doctor's/
pharmacist's advice in children 2 to 6
years of age.

Keep medicine out of reach of children/
Jauhkan ubat dari kanak-kanak

For Indication, Warning, Precaution & Side Effect.
Please refer package insert.

**MUCOLYTIC EXPECTORANT
FOR RELIEF OF COUGH**

Route of Administration: Oral

REXOM UNIZET SYRUP may cause severe
allergy and serious skin reactions. Stop using
REXOM UNIZET SYRUP and seek medical
assistance immediately if you experience any of
the following symptoms:

- 1) **Severe allergy:** breathing difficulties, light
headedness, skin swellings or rash.
- 2) **Severe skin reaction:** skin reddening,
blisters, rash, fever, sore throat or eye
irritation.

Manufactured by & Product Registration Holder:
KCK PHARMACEUTICAL INDUSTRIES SDN, BHD.
Plot 61, Bayan Lepas Industrial Park Phase 4,
Lintang Bayan Lepas 1, 11900 Bayan Lepas,
Pulau Pinang, Malaysia.

B/N:
MFG:
EXP: