

WINAMOL Drops for Infants

Paracetamol (100mg/ml)

What is in this leaflet

1. What WINAMOL Drops is used for
2. How WINAMOL Drops works
3. Before you use WINAMOL Drops
4. How to use WINAMOL Drops
5. While you are using WINAMOL Drops
6. Side effects
7. Storage and disposal
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of revision

What WINAMOL Drops is used for

For the relief of fever and mild to moderate pain. WINAMOL Drops is an analgesic in teething.

How WINAMOL Drops works

WINAMOL Drops contains Paracetamol. It works by relieving pain and reducing high temperature and fever.

Before you use WINAMOL Drops

-When you must not use it

Tell your doctor before using WINAMOL Drops if you have hypersensitivity to any of the ingredients.

-Pregnancy and breastfeeding

No ill effects due to paracetamol used in the recommended dosage, but patients should follow the advice of their doctor regarding its use. Paracetamol is excreted in breast milk but not in a clinically significant amount.

-Before you start to use it

This preparation contains PARACETAMOL. Do not take any other paracetamol containing medicines at the same time.

-Taking other medicines

Inform your doctor or pharmacist before taking this medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and WINAMOL Drops may interfere with each other. They may be affected by WINAMOL Drops or may affect how well it works.

Before taking WINAMOL Drops for Infants, make sure to consult the

physician if the patient is taking warfarin or similar medicines used to thin the blood.

How to use WINAMOL DROPS

-How much to use

Always use WINAMOL Drops exactly as your doctor has prescribed. You should check with your doctor or pharmacist if in doubt.

The usual dose:

1-3 months (4-6 kg) : 0.6-0.9 mL (60-90mg)

3-6 months (6-8 kg) : 0.9-1.2 mL (90-120mg)

6-12 months (8-10 kg) : 1.2-1.5 mL (120-150mg)

1-2 years (10-12 kg) : 1.5-1.8 mL (150-180mg)

- For an accurate dosing, weigh your child and work out the dosage to be given based on a paracetamol 15 mg for every kg body weight.
- Do not give more than 4 doses in any 24 hour period.
- Leave at least 4 hours between doses.
- Do not give this medicine to your child for more than 3 days without speaking to your doctor or pharmacist.

-When to use it

Use as directed by your doctor or pharmacist.

-How long to use it

Continue taking WINAMOL Drops as your doctor recommended. If symptoms persist, consult your doctor.

-If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

-If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of

this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Early symptoms of paracetamol overdose include pallor, nausea, vomiting, anorexia and abdominal pain. Liver damage may become apparent 12 to 48 hours after ingestion. Abnormalities of glucose metabolism and metabolic acidosis may occur.

While you are using WINAMOL Drops

Things you must do

- Inform your doctor or pharmacist if symptoms persist after taking this medication.
- Take your medicine exactly as your doctor has told you.
- Tell all the doctors, dentists and pharmacists treating you that you are taking WINAMOL Drops
- Tell your doctor immediately if you become pregnant while taking this medication.

- Things you must not do

- Do not take more than the recommended dose unless your pharmacist or doctor tells you.
- Do not take any new medicines without consulting your doctor.
- Do not give WINAMOL Drops to anyone else, even if they have the same symptoms or condition as you.

-Things to be careful of

Allergy alert: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

Consult the physician before use if the child has liver or kidney problems and if the child's symptoms do not improve. Do not exceed the stated dose.

WINAMOL Drops for Infants contains paracetamol. It should not be used with other paracetamol-containing products.

WINAMOL Drops for Infants

Paracetamol (100mg/ml)

Side effects

Like all medicines, WINAMOL Drops can cause side effects, although not everybody gets them.

If symptoms persist, do not take the medicine and refer to the doctor.

Tell your doctor if you notice any of the following side effects or notice any other effects not listed.

Cutaneous hypersensitivity reactions including skin rashes, angioedema, Stevens Johnson Syndrome/Toxic Epidermal Necrolysis have been reported.

Stop taking WINAMOL Drops for Infants and inform the physician immediately if the child: Experience allergic reactions eg, skin rash or itching, sometimes with breathing problems or swelling of the lips, tongue, throat or face; experience a skin rash or peeling, or mouth ulcers; have previously experienced breathing problems with aspirin or nonsteroidal anti-inflammatory drugs, and experience a similar reaction with WINAMOL Drops for Infants; experience unexplained bruising or bleeding, these reactions are rare.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my

[Consumer→Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and disposal of WINAMOL Drops

-Storage

Keep the bottle tightly closed protected from light. Store below 30°C.

Keep medicine out of reach of children.

-Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product description

-What it looks like

Red colour syrup with raspberry flavour.

-Ingredients

-Active ingredient
Paracetamol

-Inactive ingredients

Propylene Glycol, Polyethylene Glycol-PEG 400, Glycerin, Methyl Paraben, Propyl Paraben, Sucralose, Sodium Chloride, Raspberry Flavour, Citric Acid Anhydrous, Carmoisine Colour, Sodium Citrate and purified water.

-MAL number

MAL19086013XZ

Manufacturer and Product

Registration Holder

KCK Pharmaceutical Industries Sdn. Bhd.

Plot 61, Bayan Lepas Industrial Park
Phase 4, Lintang Bayan Lepas 1,
11900 Bayan Lepas, Pulau Pinang,
Malaysia.

Date of revision

24/07/2023

Serial Number

NPRA(R2)19/100