

Unimed's ChlorSept 5% Antiseptic Solution

25.0% v/v of Chlorhexidine gluconate 20% solution equivalent to chlorhexidine gluconate.....5% w/v

What is in this leaflet

1. What UNIMED'S CHLORSEPT 5% is used for
2. How UNIMED'S CHLORSEPT 5% works
3. Before you use UNIMED'S CHLORSEPT 5%
4. How to use UNIMED'S CHLORSEPT 5%
5. While you are using UNIMED'S CHLORSEPT 5%
6. Side effects
7. Storage and disposal
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of revision

What UNIMED'S CHLORSEPT 5% is used for

UNIMED'S CHLORSEPT 5% is used for preoperative skin disinfection and for wounds or burns.

How UNIMED'S CHLORSEPT 5% works

UNIMED'S CHLORSEPT 5% contains Chlorhexidine gluconate as its active ingredient. It has antiseptic properties which kills bacteria and inhibit the bacterial growth.

Before you use UNIMED'S CHLORSEPT 5%

-When you must not use it

Tell your doctor before using UNIMED'S CHLORSEPT 5% if you

- have hypersensitivity (itching, skin rash, redness, swelling, breathing difficulties, light headedness, rapid heart rate development)
- anaphylactic shock.
- Allergic reaction to Chlorhexidine.

-Pregnancy and breastfeeding

No effects during pregnancy and lactation are anticipated, since systemic exposure to chlorhexidine gluconate is negligible.

-Taking other medicines

Inform your doctor or pharmacist before taking this medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and UNIMED'S CHLORSEPT 5% may interfere with

each other. They may be affected by UNIMED'S CHLORSEPT 5% or may affect how well it works.

- Incompatible with soaps or other anionic agents.

How to use UNIMED'S CHLORSEPT 5%

-How much to use

Always use UNIMED'S CHLORSEPT 5% exactly as your doctor has prescribed. You should check with your doctor or pharmacist if in doubt.

The usual dose:

- **Pre-Operative Skin Disinfection:** To be used diluted with 1: 10 in 70% alcohol for preoperative skin disinfection
- **General cleansing on skin for wounds and burns:** To be used with 1: 100 in 70% alcohol for wounds or burns.

-When to use it

Use as directed by your doctor or pharmacist.

-How long to use it

Continue using UNIMED'S CHLORSEPT 5% as your doctor recommended. If symptoms persist, consult your doctor.

-If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

-If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have used too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Using too much UNIMED'S CHLORSEPT 5% may cause:

- Haemolysis (rupture of red blood cells)
- Accidental ingestion may require gastric lavage using milk, egg white, gelatine or mild soap.

While you are using UNIMED'S CHLORSEPT 5%

Things you must do

- Inform your doctor or pharmacist if symptoms persist after using this medication.
- Use the medicine exactly as your doctor has told you.
- Tell all the doctors, dentists and pharmacists treating you that you are using UNIMED'S CHLORSEPT 5%

- Things you must not do

- Do not use more than the recommended dose unless your pharmacist or doctor tells you.
- Do not take any new medicines without consulting your doctor.
- Do not give UNIMED'S CHLORSEPT 5% to anyone with possible history of an allergic reaction to Chlorhexidine-

- Things to be careful of

Stop using this product and seek immediate medical assistance if you experience rash, itching, swelling, breathing difficulties, light-headedness or rapid heartbeat.

Side effects

Like all medicines, UNIMED'S CHLORSEPT 5% can cause side effects, although not everybody gets them. If symptoms persist, do not take the medicine and refer to the doctor.

Tell your doctor if you notice any of the following side effects or notice any other effects not listed:

- Skin and subcutaneous tissue disorder. Dermatitis, pruritus, erythema, eczema, rash, skin irritation, blisters.
- Immune system disorder.
- Potentially : Serious allergic reactions or hypersensitivity (anaphylactic shock)

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.moh.gov.my
[Public → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Unimed's ChlorSept 5% Antiseptic Solution

25.0% v/v of Chlorhexidine gluconate 20% solution equivalent to chlorhexidine gluconate.....5% w/v

Storage and disposal of UNIMED'S **Serial Number**
CHLORSEPT 5% NPRA(R2)20/112

Storage

Keep container tightly closed. Protect from heat and light. Store below 30°C.

-Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product description

-What it looks like

Clear reddish pink color solution with lavender fragrance

Mode of administration

Topical

Dosage Form

Solution

Packaging

- It is filled in 120ml HDPE bottle with red HDPE screw cap.
- It is filled in 1000ml HDPE bottle with red HDPE screw cap.

-Ingredients

-Active ingredient

25.0% v/v of Chlorhexidine gluconate
20% solution equivalent to chlorhexidine gluconate.....5% w/v

-Inactive ingredients

Isopropyl Alcohol, Nonyl Phenol Ethoxylated (NP9), Fragrance (Lavender), Glucono Delta Lactone, Ponceau 4R Colour, Purified Water.

-MAL number

Manufacturer and Product

Registration Holder

KCK Pharmaceutical Industries Sdn. Bhd.
Plot 61, Bayan Lepas Industrial Park
Phase 4, Lintang Bayan Lepas 1,
11900 Bayan Lepas, Pulau Pinang,
Malaysia.

Date of revision

3/11/2020