

# Septol Antiseptic Germicide Liquid (Chloroxylenol 5.0% w/v)

Chloroxylenol (5.0% w/v)

## What is in this leaflet

1. What Septol Antiseptic is used for
2. How Septol Antiseptic works
3. Before you use Septol Antiseptic
4. How to use Septol Antiseptic
5. While you are using Septol Antiseptic
6. Side effects
7. Storage and disposal
8. Product description
9. Manufactured by and Product Registration Holder
10. Date of revision

## What Septol Antiseptic is used for

Chloroxylenol 5.0% w/v (SEPTOL) used for first aid and medical uses such as cuts, bites, abrasions, insect stings and used for personal hygiene conditions such as douching (when medically advised), dandruff, spots and pimples.

## How Septol Antiseptic works

Septol Antiseptic contains chloroxylenol as its active ingredient. It has bactericidal and antiseptic properties.

## Before you use Septol Antiseptic

### -When you must not use it

Do not use Septol Antiseptic if you are allergic to any of the ingredients listed in this leaflet.

### -Pregnancy and breastfeeding

No effects during pregnancy are anticipated.

Application of the product to the breast is not recommended during breast feeding.

### -Taking other medicines

Inform your doctor or pharmacist before using Septol Antiseptic.

## How to use Septol Antiseptic

### -How much to use

Always use Septol Antiseptic exactly as your doctor has prescribed. You should check with your doctor or pharmacist if in doubt.

### The usual dose:

### **First Aid and Medical:**

For cuts, bites, abrasions and insect stings: wash the area with one tablespoonful of SEPTOL diluted in a

250ml of water and cover with dry gauze or lint.

### **Personal Hygiene:**

For douching (when medically advised): Two teaspoonfuls of SEPTOL diluted in 1L of warm water.

For dandruff: One tablespoonful of SEPTOL diluted in 500ml of warm water. Saturate hair and scalp for 10 minutes, then shampoo.

For spots and pimples (not for eczematous condition): Bathe the affected area daily with one tablespoonful of SEPTOL diluted in 250ml of warm water.

SEPTOL Solution can be used as a household disinfectant on surfaces or in laundry.

### **Household Cleaning:**

Laundry: 4 capful of SEPTOL to 2L of water

Floors and other hard surfaces: 5 capful of SEPTOL to 2.4L of water

Lavatories, sinks (non-acrylic), drains etc.: Use undiluted SEPTOL Solution.

### -When to use it

Use as directed by your doctor or pharmacist.

### -How long to use it

Use as directed by your doctor or pharmacist.

### -If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

### - If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have accidentally ingested this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Pulmonary aspiration of chloroxylenol-based disinfectants may result in pneumonia (acute respiratory infection that affects the lungs), acute respiratory distress syndrome, and cardiorespiratory arrest. There have been reports of death by excessive consumption.

Oral ingestion may result in pharyngeal erosion (erosion in throat area), laryngeal oedema (accumulation of fluid in tissues of any part of the larynx), stomatitis (inflammation inside of the mouth), bradycardia (slow heart rate), hypotension (low blood pressure), renal failure and CNS depression.

## While you are using Septol Antiseptic

### Things you must do

- Inform your doctor or pharmacist if symptoms persist after using Septol Antiseptic
- Tell all the doctors, dentists and pharmacists treating you that you are using Septol Antiseptic.

### - Things you must not do

- Do not use more than the recommended dose unless your pharmacist or doctor tell you.

### -Things to be careful of

- Do not use on dry skin condition
- Avoid contact with the eye

If swallowed, wash out the mouth and drink plenty of water or milk.

Please consult doctor or go to the Emergency Department of your nearest hospital.

## Side effects

If you have any side effect after using this product, talk to your pharmacist or doctor.

Tell your doctor if you notice any of the following side effects or notice any other side effects:

- Hypersensitivity
- Skin sensitization
- Contact dermatitis (eczema triggered by contact with a particular substance)
- skin discolouration
- application site burn

# Septol Antiseptic Germicide Liquid (Chloroxylonol 5.0% w/v)

Chloroxylonol (5.0% w/v)

---

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website [npra.gov.my](http://npra.gov.my) [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

**Serial Number**  
NPRA(R2)22/065

## **Storage and disposal of Septol Antiseptic**

### Storage

Keep container tightly closed. Protect from heat and light. Store below 30°C.

Keep medicine out of reach of children.

### -Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

## **Product description**

### -What it looks like

A light to dark, amber colored solution with pine fragrance.

### -Ingredients

-Active ingredient  
Chloroxylonol (5.0%w/v)

-Inactive ingredients  
Potassium Hydroxide, Oleic Acid, Virgin castor oil, Terpineol, Denatured Alcohol 96% (Ethanol with Tert- butyl Alcohol), Liquid caramel and purified water.

### -MAL number

MAL

## **Manufactured by and Product**

### **Registration Holder**

KCK Pharmaceutical Industries Sdn. Bhd.  
Plot 61, Bayan Lepas Industrial Park  
Phase 4, Lintang Bayan Lepas 1,  
11900 Bayan Lepas, Pulau Pinang,  
Malaysia.

### **Date of revision**

24/08/2022