
UniMoist Aqueous Cream BP

Composition and Strength of Active ingredients:

Anionic Emulsifying Wax BP	9.0% w/w
Liquid Paraffin BP.....	6.0% w/w
White Soft Paraffin BP.....	15.0% w/w

Preservatives:

Methyl Paraben	0.15% w/w
Propyl Paraben	0.08% w/w

Product Description

White to off-white colour, smooth, homogenous cream.

Pharmacodynamics

White soft paraffin and liquid paraffin have emollient properties.

Pharmacokinetics

None applicable.

Indications

For topical application to the skin as an emollient for the symptomatic relief of dry skin conditions and as soap-substitute for skin washing.

Recommended Dose

Apply a thin layer over affected area 2 to 3 times daily or as required.

Route of Administration

Topical

Contraindications

Hypersensitivity to any of the ingredients.

Warning and Precautions

For external use only.

Aqueous cream contains sodium lauryl sulfate which may cause local skin reactions (e.g. stinging and contact dermatitis), particularly in children with atopic eczema.

Also contains cetostearyl alcohol which may cause local skin reactions (e.g. contact dermatitis).

Do not smoke or go near naked flames due to risk of severe burns. Fabric (clothing, bedding, dressings etc) that has been in contact with this product burns more easily and is a serious fire hazard. Washing clothing and bedding may reduce product build-up but not totally remove it.

Keep all medicines out of the sight and reach of children.

Interactions with Other Medicaments

No clinically significant drug interactions known.

Pregnancy & Lactation

The safety of aqueous cream during pregnancy and lactation has not been established but is not considered to constitute a hazard during these periods.

Side Effects

Occasional allergic reactions.

Aqueous cream may be associated with immediate cutaneous reactions when used as a leave-on emollient, such as stinging, burning, itching and redness.

Symptoms and Treatment of Overdose

Symptoms of accidental oral ingestion may include gastrointestinal symptoms, particularly diarrhoea. Treatment need only be symptomatic.

Effects on Ability to Drive and Use Machine

No adverse effects known.

Storage Condition

Keep container tightly closed. Store below 30°C. Protect from freezing, heat and direct light.

Dosage form and packaging available

100gm & 500gm are packed into HDPE jar.

Registration Number:

MALXXXX

Manufactured by:**KCK Pharmaceutical Industries Sdn. Bhd.**

No. 5 & 7 Hala Beringin, Diamond Valley Industrial Park,
11960 Bayan Lepas, Pulau Pinang, Malaysia.

Product Registration Holder:**KCK Pharmaceutical Industries Sdn. Bhd.**

Plot 61, Bayan Lepas Industrial Park Phase 4,
Lintang Bayan Lepas 1, 11900 Bayan Lepas,
Pulau Pinang, Malaysia

Date of Revision:

30 June 2025