



biosyn Arzneimittel GmbH
Schorndorfer Strasse 32
70734 Fellbach, Germany

Product Information

1. NAME OF THE MEDICINAL PRODUCT

selenase® 100 micrograms,
solution for injection (50 micrograms/ml)

selenase® 500 micrograms,
solution for injection (50 micrograms/ml)

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

selenase® 100 micrograms, solution for injection (50 micrograms/ml)
Each 2 ml injection ampoule contains 100 micrograms selenium in the form of 333 micrograms sodium selenite pentahydrate ($\text{Na}_2\text{SeO}_3 \cdot 5 \text{H}_2\text{O}$), corresponding to 50 µg/ml.

selenase® 500 micrograms, solution for injection (50 micrograms/ml)
Each 10 ml injection vial contains 500 micrograms selenium in the form of 1.66 mg sodium selenite pentahydrate ($\text{Na}_2\text{SeO}_3 \cdot 5 \text{H}_2\text{O}$), corresponding to 50 µg/ml.

For excipients, see 6.1

3. PHARMACEUTICAL FORM

Solution for injection.
Clear and colourless solution.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Treatment of proven selenium deficiency that cannot be offset from food sources.

4.2 Posology and methods of administration

Daily dose:

100–200 micrograms selenium (equivalent to 1–2 ampoules). If more selenium is necessary to reach the normal blood level, this dose can be increased to 500 micrograms selenium (equivalent to 5 ampoules = 5 x 100 micrograms or 1 injection vial = 500 micrograms, respectively).

Method of administration

selenase® solution for injection is administered as an intramuscular or intravenous injection. Selenium levels in whole blood or serum should be determined in order to monitor the success of therapy.

When selenase® solution for injection is administered as a supplement to general infusion solutions for total parenteral nutrition, a daily dose of 100 micrograms selenium must be ensured.

There is no time limit to the administration of selenase® solution for injection in a supplementary dose (100 micrograms selenium per day, equivalent to 2 ml selenase® solution for injection).

Do not take selenium together with or too close to strong reducing substances as vitamin C. Vitamin C (Ascorbic acid) has an extremely high redox potential and reduces selenite to elementary selenium which is not harmful, but also not bioavailable. Because food may contain also strong reducing substances, please leave an interval of at least 1 hour between eating and drinking (of fruit juices, not normal water) and the intake of selenase®.

Dosage in children and in special patient groups

No scientific evidence exists which would require dosage adjustment in children or patients with renal or hepatic impairment.

For children the dosage recommendation is 2 mcg/kg body weight. But it may be necessary to give higher dosages to a maximum of 30 micrograms daily for children with body weight lesser than 40 kg according to the determination of selenium concentration in whole blood. Children with body weight higher than 40 kg may be treated with dosages for adults.

Dosage in patients with renal or hepatic impairment

There is no scientific evidence on dosage adjustment in patients with renal or hepatic impairment.

4.3 Contraindications

Hypersensitivity to any component of the product (active substance or excipients).
Selenosis.

4.4 Special warning and precautions for use

selenase® solution for injection contains less than 1 mmol sodium (23 mg) per ml.

4.5 Interaction with other medicinal products and other forms of interaction

When preparing an infusion solution with selenase® solution for injection as a supplement, it must be ensured that the pH value does not fall below 7.0 and that the solution is not mixed with reducing substances (e.g. vitamin C), as it may result in precipitation of elemental selenium (see section 6.2 "Incompatibilities"). Elemental selenium is not soluble in an aqueous medium and has no biological availability.

4.6 Pregnancy and lactation

Pregnancy:

There are no data from the use of selenase® in pregnant woman. Limited published data from animal studies reveal only evidence for toxicity to reproduction at maternally toxic dose.

No adverse effect of sodium selenite on the pregnancy or unborn child is expected, provided that it is used in case of proven selenium deficiency.

Lactation:

Selenium is excreted in breast milk. Doses correcting selenium deficiency in breast feeding woman are not expected to exert adverse effects on the suckling infant.

4.7 Effects on ability to drive and use machines

None

4.8 Undesirable effects

No adverse effects associated with the use of selenase are to be expected if selenase is used according to the manufacturer's product recommendations.

However, there have been isolated reports/ individual cases of the following adverse events:

Severe nausea, abdominal cramping, diarrhoea, clouding of consciousness after selenase® solution for injection, discharge from right breast and navel two days after first administration of selenase®, rise in GOT, GPT, γ-GT, AP, abdominal pain, long lasting pyrosis, salty taste in the mouth, cardiac dysrhythmia, allergic reaction (inflammable swelling, formation of oedemas in the region of the forearm), nausea, allergy, flu-like symptom, diagnosis of tertiary tumour, exanthema, pruritus, increase blood pressure, pulse interruption time.

Note, the causal relationship of the above adverse events has not/cannot be established.

4.9 Overdose

Signs of an acute overdose are an odour of garlic on the breath, tiredness, nausea, diarrhoea and abdominal pain. Chronic overdose can affect growth of nails and hair and may lead to peripheral polyneuropathy.

Countermeasures include gastric lavage, forced diuresis or the administration of high doses of vitamin C. In the case of an extreme overdose (1,000–10,000 times the normal dose) an attempt should be made to eliminate the selenium by dialysis. Administration of dimercaprol is not recommended as the toxic effect of selenium is potentiated.

(see reverse)

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Mineral supplement
ATC code: A12C E02

Selenium is a co-factor in various enzymes in the human body and therefore belongs to the essential trace elements. To date, more than 25 proteins and protein subunits containing selenium have been identified and most clinical and biochemical effects of selenium can be attributed to their activity. However, not all the effects of selenium are exclusively related to the action of different enzymes.

Selenium-containing glutathione peroxidase and selenium protein P have been identified in humans. Glutathione peroxidase is part of the anti-oxidant protection mechanism of the cell in mammals. As a constituent of glutathione peroxidase, selenium can delay the lipid peroxidation rate and thus the resultant damage to the cell wall. Glutathione peroxidase affects the metabolism of leukotrienes, thromboxanes and prostacyclines. In animals, type I iodothyronine-5'-deiodinase is characterised as a selenium enzyme that converts thyroxine (T_4) into triiodothyronine (T_3), the active thyroid hormone.

A selenium deficiency is manifested in reduced selenium levels in whole blood or plasma and in the suppression of glutathione peroxidase activity in whole blood, plasma or thrombocytes. The pathophysiological relevance of selenium-dependent reactions has been demonstrated in studies of selenium deficiencies in humans and animals: Selenium deficiency activates and inhibits the response of immunological mechanisms, particularly non-specific cell and body fluid responses. Selenium deficiency affects the activity of various hepatic enzymes. Selenium deficiency potentiates damage occasioned to the liver by oxidative or chemical factors and the toxicity of heavy metals such as mercury and cadmium.

For humans, the following diseases are described as a consequence of selenium deficiency: Keshan disease, an endemic cardiopathy, and Kaschin-Beck disease, an endemic osteoarthropathy that is associated with very severe deformity of the joints. Clinically manifest selenium deficiency is also observed as a consequence of long-term parenteral nutrition and unbalanced diets.

5.2 Pharmacokinetic properties

Sodium selenite is not immediately converted to proteins. In the blood, the majority of the supply of selenium is used by the erythrocytes and converted to hydrogen selenide under the action of enzymes. Hydrogen selenide acts as a central pool of selenium for both elimination and the specific integration of selenium in selenoproteins. Reduced selenium binds to plasma proteins that migrate to the liver and other organs. Secondary plasma transport from the liver to the target tissues, that produce glutathione peroxidase by synthesis, probably occurs via a P-selenoprotein containing selenocysteine. The subsequent metabolic pathway of selenoprotein synthesis has to date

only been studied in prokaryotes. In the metabolic process, selenocysteine is specifically incorporated in the peptide chains of glutathione peroxidase.

All excess hydrogen selenide is metabolised via methylselenol and dimethylselenide to the trimethylselenonium ion, the principal elimination product.

After oral administration, selenium is principally absorbed from the small intestine. Absorption of sodium selenite in the intestine is not regulated by homeostatic mechanisms. Depending on the concentration of sodium selenite and the presence of related substances, it is usually between 44% and 89%, and sometimes more than 90%. The amino acid cysteine increases the absorption of sodium selenite.

The total quantity of selenium present in the human body is between 4 mg and 20 mg. Humans excrete selenium in the faeces, via the kidneys and through the respiratory system, depending on the amount administered. Selenium is predominantly eliminated in the form of the trimethylselenonium ion via the kidneys. Elimination is dependent on the selenium status.

After intravenous or oral administration, the process of selenium elimination was divided into three phases. After oral administration of 10 micrograms in the form of [^{75}Se] sodium selenite, 14–20% of the absorbed selenium is eliminated via the kidneys in the first two weeks, while almost nothing was eliminated via the lungs and skin. The retention of selenium in the whole body decreased in three phases, with half-lives of 0.7–1.2 days in phase 1, 7–11 days in phase 2 and 96–144 days in phase three. The selenium concentration decreased more rapidly in the liver, heart and plasma than in the skeletal muscles or in the bones. Of an intravenously administered dose of [^{75}Se] sodium selenite, 12% was excreted in the first 24 hours. A further 40% was eliminated with a biological half-life of 20 days. The half-life of the third phase was 115 days.

Elimination after oral and intravenous administration of a physiological dose of [^{74}Se] sodium selenite was compared directly: after administration of 82 micrograms selenium in the form of sodium selenite, 18% of the intravenous dose and 12% of the oral dose was eliminated via the kidneys in the first 24 hours together with metabolised physiological selenium. After this phase, the process of elimination by both routes of administration is more or less the same. In healthy volunteers, the elimination of orally and parenterally administered sodium selenite was comparable.

5.3 Preclinical safety data

Published literature on single and repeated dose toxicity, genotoxicity and carcinogenicity of selenium and sodium selenite reveals no evidence for adverse health effects in addition to those already known from experience in humans. Toxicity to reproduction was only found at maternally toxic doses.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Sodium chloride, hydrochloric acid, water for injections.

6.2 Incompatibilities

When preparing an infusion solution with selenase[®] solution for injection as a supplement, it must be ensured that the pH value does not fall below 7.0 and that the solution is not mixed with reducing substances (e.g. vitamin C), as it may result in precipitation of elemental selenium. On grounds of safety, non-specific precipitation should be avoided after mixing infusion solutions with selenase[®] solution for injection.

6.3 Shelf-life

Four (4) years from the date of manufacture. Use immediately after opening.

6.4 Special precautions for storage

Do not store above 30 °C.

6.5 Nature and contents of container

selenase[®] 100 micrograms, solution for injection (50 micrograms/ml)
Injection ampoules each containing 2 ml of injection solution are made of glass. Pack size: 10

selenase[®] 500 micrograms, solution for injection (50 micrograms/ml)
Injection vials each containing 10 ml of solution for injection are made of glass with a rubber stopper.
Pack size: 10

6.6 Instructions for use and handling

For single use only. Discard any unused contents. selenase[®] may be mixed with 0.9% NaCl.

7. MANUFACTURER

selenase[®] 100 micrograms, solution for injection (50 micrograms/ml)

Manufactured by:
Solupharm Pharmazeutische Erzeugnisse GmbH
Industriestrasse 3, 34212 Melsungen, Germany

selenase[®] 500 micrograms, solution for injection (50 micrograms/ml)

Manufactured by:
Wasserburger Arzneimittelwerk GmbH
Herderstrasse 2, 83512 Wasserburg, Germany

Released by:
biosyn Arzneimittel GmbH
Schorndorfer Str. 32, 70734 Fellbach, Germany

8. DATE OF REVISION OF THE TEXT

07 July 2025