

**XIDINESOL 4% w/v**

VIXIDXX-M

**Description**

Red viscous solution with mild floral fragrance.

**Composition**

Chlorhexidine Gluconate 4%

**Indications**

Antiseptics & Disinfectants

**Contraindications**

Xidinesol preparations are contraindicated for patients who have previously shown a hypersensitivity reaction to chlorhexidine. However, such reactions are extremely rare.

**Precautions**

Since chlorhexidine is irritant, it is recommended that it should not be used on the brain, meninges, middle ear, or other sensitive tissues.

**Interaction with Other Medicaments and Other Forms of Interaction**

Chlorhexidine is incompatible with soap and other anionic agents. Hypochlorite bleaches may cause brown stains to develop in fabrics which have previously been in contact with preparations containing chlorhexidine.

**Main Side/Adverse Effects**

Skin sensitivity to chlorhexidine has occasionally been reported. Severe hypersensitivity reactions, including anaphylactic shock, have been reported rarely after topical use of chlorhexidine.

**Overdose**

This has not been reported.

Accidental ingestion: Chlorhexidine taken orally is poorly absorbed. Treat with gastric lavage using milk, raw egg, gelatin or mild soap. Employ supportive measures as appropriate.

**Use Instruction**

For external use only. Do not dilute. To be used full strength.

Storage : Store below 30°C.  
Presentation/Packing : Bottle of 800 ml, 850 ml, 900 ml, 1000 ml.  
Manufactured by : HOVID Bhd., 121, Jalan Tunku Abdul Rahman, 30010 Ipoh, Malaysia.  
Revision date : June 2011