

DESCRIPTION

White cream with sheen.

PHARMACODYNAMICS

Miconazole is an imidazole antifungal agent; it is fungicidal to various species of Trichophyton, Epidermophyton, Microsporum, Candida, Cryptococcus, Aspergillus, Coccidioides, Paracoccidioides, Cladosporium, Phialophora, Madurella, Blastomyces and Histoplasma.

The drug permeates the chitin of the fungal wall and increases the membrane permeability to various intracellular substances. Miconazole is also effective against gram-positive bacteria. The drug readily penetrates the stratum corneum of the skin and persists there for more than 4 days after application.

In the treatment of tinea pedis, topical miconazole relieves itching within a few days; vesicles and fissures heal rapidly but desquamation may continue for several weeks. In the treatment of tinea cruris, the drug is comparably effective. Tinea versicolor, ringworm, onychomycosis and cutaneous candidiasis also respond to topical miconazole.

PHARMACOKINETICS

Absorption

Very little is absorbed systemically following topical application.

Excretion

Only trace amounts can be found in the urine.

INDICATIONS

For the topical treatment of tinea pedis, tinea cruris and tinea corporis, and in topical treatment of cutaneous candidiasis, tinea and pityriasis versicolor.

WARNINGS AND PRECAUTIONS

- In patients on warfarin, caution should be exercised and the anticoagulant effect should be monitored (see Interactions).
- Do not apply occlusive dressing over this medication unless directed to do so by physician.
- Avoid contact with the eyes.
- If irritation occurs or if there is no improvement within 4 weeks, discontinue use and consult a doctor or pharmacist.
- Do not use in children under 2 years of age except under the advice and supervision of a doctor.

MAIN SIDE/ADVERSE EFFECTS

Local irritation and occasionally sensitisation may occur at the site of application.

OVERDOSAGE

Clinical features: Ingestion-mucous membrane irritation, anorexia, nausea, vomiting, diarrhoea.

Treat poisoning by the necessary symptomatic measures. Gastric lavage is probably unnecessary.

Note: The prescribed course should be completed. The information given here is limited. For further information consult your doctor or pharmacist.

hovid

500 Grams

Store below 30°C.
Keep out of reach of children. *Jauhi daripada kanak-kanak.*

REG. NO. : MAL 19900252AZ
SIN06466P
HK-35640
BRU16032014P

TINAZOL CREAM 2%

MICONAZOLE NITRATE CREAM B.P.

Contents:

Miconazole Nitrate 2.0 % w/w

Preservative:

Chlorocresol 0.2 % w/w

Indications and direction (Adults and children):

- Tinea pedis, tinea cruris, tinea corporis or cutaneous candidiasis - topical to the skin, one or two times a day for 2 to 4 weeks.
- Tinea versicolor - topical to the skin, once a day

FOR EXTERNAL USE ONLY

PRODUCT REGISTRATION HOLDER /
MANUFACTURED BY: **HOVID Berhad**
121, JALAN TUNKU ABDUL RAHMAN
(JALAN KUALA KANGSAR),
30010 IPOH, PERAK, MALAYSIA.

Controlled
Medicine

Lot :

Mfg :

Exp :

LTIN05



VL TIN05-var2