

# FLUCOR NIGHT SOFT GELATIN CAPSULE

Paracetamol / Pseudoephedrine Hydrochloride / Chlorpheniramine Maleate (250mg / 30mg / 2mg)

## What is in this leaflet

1. What Flucor Night is used for
2. How Flucor Night works
3. Before you use Flucor Night
4. How to use Flucor Night
5. While you are using Flucor Night
6. Side effects
7. Storage and Disposal of Flucor Night
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

## What Flucor Night is used for

Flucor Night is used for the temporary relief of running nose and sneezing, minor aches and pain, headache, fever and nasal congestion associated with the common cold.

## How Flucor Night works

This product contains paracetamol, pseudoephedrine hydrochloride and chlorpheniramine maleate as active ingredients. Paracetamol works by stopping the pain message from getting through the brain. Paracetamol also reduces fever. Pseudoephedrine hydrochloride is a decongestant which unblocks your nose and sinuses to help you breathe easier. Chlorpheniramine maleate is an antihistamine which helps to relieve the symptoms of allergies.

## Before you use Flucor Night

### - When you must not use it

Do not use Flucor Night if you or your children are:

- Allergic to paracetamol, pseudoephedrine, chlorpheniramine, other antihistamines or any of the other ingredients in the product (listed in Product Description).
- Having severe high blood pressure or severe heart disease

- Taking or have taken medications that treat depression (monoamine oxidase inhibitors / MAOIs) in the past two weeks

### - Before you start to use it

Tell your doctor if you or your children:

- Have impaired kidney or liver function
- Have breathing difficulties, persistent cough or glaucoma (high pressure in the eye)
- Have heart diseases, thyroid diseases, diabetes or high blood pressure
- Have difficulty in urination due to enlargement of the prostate gland
- Are pregnant, planning for pregnancy or breast-feeding before using any medicine.

### - Taking other medicines

Tell your doctor or pharmacist before taking Flucor Night if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop. Medication that may interact with Flucor Night includes:

- Decongestants
- Certain medications that treat depression
- Medications that suppress appetite
- Certain medications that treat attention deficit hyperactivity disorder (ADHD)
- Medications that reduce blood pressure such as guanethidine, debrisoquine, and methyl dopa.

## How to use Flucor Night

For oral administration only.

Follow all directions given to you by your doctor and pharmacist

carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

### - How much to use

Adults and children 12 years of age or over: Take 2 capsules every 4 hours; not exceeding 8 capsules in 24 hours.

### - When to use it

Use 4 times daily or as directed by your doctor or pharmacist. Minimum dosing interval: 4 hours.

### - How long to use it

Take Flucor Night for the duration as your doctor recommends.

### - If you forget to use it

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

### - If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have used too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too much Flucor Night may cause vomiting, stomach pain, low blood pressure and sweating.

# FLUCOR NIGHT SOFT GELATIN CAPSULE

Paracetamol / Pseudoephedrine Hydrochloride / Chlorpheniramine Maleate (250mg / 30mg / 2mg)

## While you are using Flucor Night

### - Things you must do

- Tell all the doctors, dentists and pharmacists treating you that you are taking Flucor Night.
- Check with your doctor if symptoms do not improve or recurs.
- Long term use should be avoided unless directed by doctor
- Tell your doctor immediately if you become pregnant while taking this medication.

### - Things you must not do

- **WARNING:** This preparation contains PARACETAMOL. Do not take with other paracetamol-containing medicines at the same time.
- Do not use any new medicines without consulting your doctor or pharmacist.
- Do not give Flucor Night to anyone else, even if they have the same symptoms or condition as you.

### - Things to be careful of

Avoid drinking alcohol with this medicine.

### Driving and using machines

This medicine may affect your ability to drive or use machines. If the tablet makes you drowsy or tired, do not drive or use machines.

## Side effects

**ALLERGY ALERT:** Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters and rash. These

could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance immediately.

This product contains sodium metabisulphite that may cause serious allergic reactions in certain people.

When using this product you also may experience:

- Blood disorders such as low white or red blood cell count
- Increased heart rate, anxiety, restlessness, difficulty to sleep and to pass urine
- Painful urination, drowsiness, increased sweating and loss of appetite.

Visit your doctor or pharmacist if you experience any side effects after using this medicine. You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction monitoring visiting the website [npra.gov.my](http://npra.gov.my) [Consumers → Reporting Side Effects to Medicines (ConSERF) or vaccines (AEFI)]

## Storage and Disposal of Flucor Night

### - Storage

Store Flucor Night below 25°C. Protect from light. Keep out of reach and sight of children.

### - Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

## Product Description

### - What it looks like

Oblong, dark green soft gelatin capsule printed with 'Flucor', filled with transparent green liquid.

### - Ingredients

#### - Active ingredients

- Paracetamol, pseudoephedrine hydrochloride, chlorpheniramine maleate.

#### - Inactive ingredients

- Quinoline Yellow Extra, FD&C Blue No.1, Polysorbate 80, Sodium Metabisulphite, Glycerin, Polyvinylpyrrolidone K-25, Deionised Water, Sodium Acetate Trihydrate, Propylene Glycol, Polyethylene Glycol 400, Gelatine, Sorbital Solution 70%

### - MAL number(s)

MAL20041202AZ

## Manufacturer and Product Registration Holder

HOVID Bhd.  
121, Jalan Tunku Abdul Rahman  
(Jalan Kuala Kangsar), 30010 Ipoh,  
Perak, Malaysia

## Date of revision

06/03/2024

## Serial Number

NPRA(R4/1)050324/00027