

DISUF-B CREAM

Fusidic Acid / Betamethasone valerate (2.0% w/w / 0.1% w/w)

What is in this leaflet

1. What Disuf-B Cream is used for
2. How Disuf-B Cream works
3. Before you use Disuf-B Cream
4. How to use Disuf-B Cream
5. While you are using it
6. Side effects
7. Storage and Disposal of Disuf-B Cream
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Disuf-B Cream is used for

Disuf-B Cream is used to treat bacterial skin infection and inflammation.

How Disuf-B Cream works

Disuf-B Cream contains two active ingredients which are fusidic acid, an antibiotic and betamethasone valerate from steroid group.

Fusidic acid works by killing the bacteria that cause infections and betamethasone help to reduce any swelling, redness or itchiness of your skin.

Before you use Disuf-B Cream

- When you must not use it

Do not use this medicine if:

- You are allergic (hypersensitive) to fusidic acid or betamethasone valerate or any other ingredients of this medicine (listed in Product Description).
- You have red spotty rash around your mouth or chin.

- You have viral or fungal skin infection.
- You have acne-like breakouts on face.
- You have tuberculosis of the skin (skin infection).
- You have skin ulceration.

- Before you start to use it Pregnancy and lactation

- Please consult your doctor or pharmacist if you are pregnant, planning for pregnancy or breast-feeding before using any medicine.
- Disuf-B must only be used during pregnancy if the benefits is more than the risks to the fetus. However, topical steroid should not be used extensively in pregnancy.

- Taking other medicines

- Tell your doctor or pharmacist if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

How to use Disuf-B Cream

- How much to use

- Wash your hands before and after using this cream
- Apply sparingly to the affected area.
- Follow all directions given to you by your doctor and pharmacist carefully.

- When to use it

- Uncovered lesions: 2-3 times daily.
- Covered lesions: Less frequent application may be adequate.

- Use as directed by your doctor or pharmacist.

- How long to use it

- Use Disuf-B Cream for the duration that the doctor or pharmacist recommends.
- Do not use more than 7 days. Inform your doctor if your condition does not improve or is worsening.

- If you forget to use it

Apply the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to apply the medicine and skip the missed dose. Do not apply double dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you or anyone else may have using too much of this medicine or accidentally ingested this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using it

- Things you must do

- Use your medicine exactly as your doctor or pharmacist has told you.
- Tell your doctor or pharmacist treating you that you are using Disuf-B Cream.
- Tell your doctor immediately if you become pregnant while using this medicine.

DISUF-B CREAM

Fusidic Acid / Betamethasone valerate (2.0% w/w / 0.1% w/w)

- Things you must not do

- Do not stop using the medicine unless advised by your doctor.
- Do not take any new medicines without consulting your doctor or pharmacist.
- Do not give Disuf-B Cream to anyone else, even if they have the same symptoms or conditions as you.

- Things to be careful of

- This medicine is for external use only.
- If you are using this medicine on your face or near eyes, avoid contact with eyes as it may cause glaucoma (increase pressure in eyes).
- Consult doctor or pharmacist before using on extensive surface or under occlusive dressings including baby napkins in infants and children.

Side effects

Like all medicines, Disuf-B Cream can cause side effects, although not everybody gets them.

You must get urgent medical help if you have any of the following symptoms of hypersensitivity:

- Difficulty breathing
- Face or throat swelling
- Skin rash, redness or itchiness.

- Other possible side effects:

- Thinning of the skin
- Stretch marks
- Adrenal insufficiency; symptoms include tiredness, low blood pressure and blood sugar, and gastrointestinal disturbances.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of Disuf-B Cream

- Storage

- Keep out of the reach and sight of children.
- Do not use Disuf-B Cream after the expiry date on the carton.
- Store below 25°C.

- Disposal

Medicines should not be thrown away in waste water or in household waste. Please ask your pharmacist how to throw away any medicine you do not need any more. If you do this you will help protect the environment.

Product Description

- What it looks like

White, opaque cream.

- Ingredients

- Active ingredient(s):
Fusidic acid 2.0% w/w and Betamethasone valerate 0.1% w/w
- Inactive ingredients:
Propyl hydroxybenzoate, butylated hydroxytoluene, methyl hydroxybenzoate, aminomethyl propanol, polysorbate 80, carbomer (Carbopol 934), liquid paraffin, propylene glycol, sorbitan monoleate and purified water.

- MAL number:

MAL08111773AZ

Manufacturer

HOVID Bhd.
121, Jalan Tunku Abdul Rahman (Jalan Kuala Kangsar), 30010 Ipoh, Perak, Malaysia.

Product Registration Holder

HOVID Bhd.
121, Jalan Tunku Abdul Rahman (Jalan Kuala Kangsar), 30010 Ipoh, Perak, Malaysia.

Date of revision

16/03/2023

Serial Number

NPRA (R4/1)280223/00019