

HOVID FEXOVID TABLET® Consumer Medication Information Leaflet (RiMUP)

Fexofenadine hydrochloride (180mg)

What is in this leaflet

1. What Hovid Fexovid Tablet is used for
2. How Hovid Fexovid Tablet works
3. Before you use Hovid Fexovid Tablet
4. How to use Hovid Fexovid Tablet
5. While you are using it
6. Side effects
7. Storage and Disposal of Hovid Fexovid Tablet
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Hovid Fexovid Tablet is used for

Hovid Fexovid Tablet is used to treat allergic symptoms like sneezing, runny nose, itchy eyes, nose, and throat, as well as red and watery eyes.

It also treats hives of unknown cause and reduce itchiness.

How Hovid Fexovid Tablet works

Hovid Fexovid Tablet contains fexofenadine hydrochloride, which is an antihistamine. It works by blocking the action of histamine in order to treat allergy symptoms.

Before you use Hovid Fexovid Tablet

- When you must not use it

Do not take this medicine if:

- You are allergic to fexofenadine hydrochloride or any other ingredients of this medicine (listed in Product Description).
- You have kidney problem.

- Before you start to use it

Tell your doctor if :

- You have liver problem.
- You have heart problem.

If you are unsure of anything, it is essential that you ask your doctor or pharmacist for advice.

Pregnancy and lactation

Please consult your doctor or pharmacist if you are pregnant, planning for pregnancy or breast-feeding before using any medicine.

- Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket, or health food shop.

Tell your doctor if you are currently taking:

- Erythromycin (antibiotic)
- Ketoconazole (antifungal)
- Antacids, including aluminum and magnesium hydroxide-containing (antiulcer medications)

How to use Hovid Fexovid Tablet

Shallow your tablet orally in whole with plenty of water.

- How much to use

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

Adult and children 12 years old and above :

Take 1 tablet, once per day.

- When to use it

Take once per day or as directed by your doctor or pharmacist.

- How long to use it

Take Hovid Fexovid Tablet for the duration that the doctor recommends.

- If you forget to use it

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you or anyone else may have use too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too many tablets may cause dizziness, drowsiness, and dry mouth.

While you are using it

- Things you must do

Tell all the doctors, dentists and pharmacists treating you that you are using Hovid Fexovid Tablet.

Tell your doctor immediately if you become pregnant while using this medication.

- Things you must not do

Do not stop using the medicine unless advised by your doctor.

Do not use any new medicines without consulting your doctor or pharmacist.

Do not give Hovid Fexovid Tablet to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

Driving and using machines

This medicine may affect your ability to drive or use machines. If the tablets make you feel sick, dizzy or tired, or give you a headache, do not drive or use machines and contact your doctor immediately.

Side effects

Like all medicines, Hovid Fexovid Tablet can cause side effects, although not everybody gets them.

The side effects include :

- Back pain
- Coughing
- Dizziness
- Drowsiness
- Menstrual pain

HOVID FEXOVID TABLET® *Consumer Medication Information Leaflet (RiMUP)*

Fexofenadine hydrochloride (180mg)

- Stomach upset
- Fatigue
- Headache
- Nausea
- Inflammation or infection in the middle ear
- Inflammation of the sinus
- Cold and flu

Talc
Hydroxypropyl methylcellulose E-5
Hydroxypropyl methylcellulose E-15
Titanium dioxide
Iron oxide red
Sunset yellow lake
Polyvinyl pyrrolidone
Magnesium stearate
Croscarmellose sodium

Visit your doctor or pharmacist immediately if you experience any side effects after using this medicine.

- MAL number(s):
MAL11040011AZ

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring visiting the website npra.gov.my [Consumers→ Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Manufacturer

HOVID Bhd
Lot 56442, 7 ½ Miles,
Jalan Ipoh/Chemor,
31200 Chemor,
Perak, Malaysia.

Product Registration Holder

HOVID Bhd
121, Jalan Tunku Abdul Rahman
(Jalan Kuala Kangsar),
30010 Ipoh,
Perak, Malaysia

Storage and Disposal of Hovid Fexovid Tablet

- Storage
Keep out of the reach and sight of children.

Store below 30°C.

- Disposal
Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Date of revision

27/03/2024

Serial Number

NPRA(R4/1)260324/00048

Product Description

- What it looks like
Oblong, peach film-coated tablet plain on both sides with bevel-edged and shallow convex faces.
- Ingredients
 - Active ingredient
Fexofenadine hydrochloride
 - Inactive ingredients
Microcrystalline cellulose
Pregelatinized starch
Isopropyl alcohol
Propylene glycol