

HOSOLVON TABLET

Bromhexine Hydrochloride (8mg)

What is in this leaflet

1. What Hosolvon Tablet is used for
2. How Hosolvon Tablet works
3. Before you use Hosolvon Tablet
4. How to use Hosolvon Tablet
5. While you are using Hosolvon Tablet
6. Side effects
7. Storage and disposal of Hosolvon Tablet
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Hosolvon Tablet is used for

Hosolvon Tablet is used in conditions where there are a lot of thick secretions or mucus in your air passages.

How Hosolvon Tablet works

Hosolvon Tablet helps to clear the chest by thinning the mucus in your air passages.

Before you use Hosolvon Tablet

- When you must not to use it

Do not use Hosolvon Tablet if you are allergic to bromhexine or to any of the other ingredients in Hosolvon Tablet.

-Before you start use it

Before taking Hosolvon Tablet, you must tell your doctor or pharmacist if you have stomach ulcers.

If you are not sure if you have, or have had, any of these conditions, you should

raise those concerns with your doctor or pharmacist.

-Taking other medicines

You should always tell your doctor if you are taking or have recently taken any other medicines including those obtained without a prescription.

How to use Hosolvon Tablet

-How much to use

Always take Hosolvon Tablet exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.

The recommended doses are stated below:

Adults: 1 to 2 tablets 3 times daily.

Children 6 - 12 years: Half tablet 3 times daily.

Doctors sometimes prescribe different doses to these.

The label on the pack will tell you which dose you should take. If you are still not sure, ask your doctor or pharmacist.

-When to use it

Take three times daily and try to take Hosolvon Tablet at the same time each day.

-How long to use it

If your condition get worsen or you do not start to feel

better within a few days, go back and see your doctor.

-If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

-If you take too much (overdose)

If you take more Hosolvon Tablet than you should, tell a doctor or go to a hospital straight away. Take the medicine pack with you. This is so the doctor knows what you have taken. If you take too much Bromhexine, it is likely that you will get the following conditions:

- Eating disorder
- Nausea
- Vomitting
- Abdominal discomfort
- Diarrhea
- Excess phlegm
- Shallow breathing
- Seizure

While you are using Hosolvon Tablet

-Things you must do

Talk to your doctor before taking this medicine if:

- You are pregnant, might become pregnant or think you may be pregnant. This medicine should not be

HOSOLVON TABLET

Bromhexine Hydrochloride (8mg)

taken during the first three months of your pregnancy.

- You are breast-feeding or planning to breast-feed.

-Things you must not do

Do not give Hosolvon Tablet to anyone else, even if they have the same condition as you.

Do not take Hosolvon Tablet to treat any other complaints unless your pharmacist or doctor tells you to.

Do not take more than the recommended dose unless your pharmacist or doctor tells you to.

-Things to be careful of

Special care is recommended during pregnancy. The benefits of Hosolvon Tablet must be assessed against possible effects on your child.

Side Effects

Hosolvon Tablet may cause severe allergy and serious skin reactions. Stop using Hosolvon Tablet and seek medical assistance immediately if you experience any of the following symptoms:

- 1) severe allergy: breathing difficulties, light headedness, skin swellings or rash.
- 2) severe skin reaction: skin reddening, blisters, rash, fever, sore throat or eye irritation.

You should be aware that all medicines carry some risks and

that all possible risks may not be known at this stage.

Ask for the advice of your doctor or pharmacist if you have gastrointestinal disturbances or any concerns about the effects of taking Hosolvon Tablet.

Tell your doctor or pharmacist as soon as possible if you experience any side effects during or after taking Bromhexine, so that these may be properly treated. In addition, unexpected effects, not listed above, can occur with any medicine. When you take Bromhexine, you may notice an increase in mucus from your air passages. This is common when you take Hosolvon Tablet and is due to the way Bromhexine works.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npa.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and disposal of Hosolvon Tablet

-Storage

Keep all medicines out of sight and reach of children. Do not take this medicine after the expiry date stamped on the pack. Store below 30°C. Protect from light and moisture.

-Disposal

Medicines should not be disposed of via wastewater or

household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product description

-What it looks like

Round, 7.0 mm in diameter, white to off-white uncoated tablet, shallow convex faces, "HD" embossed and scored on the same face.

-Ingredients:

-Active ingredient(s):-
Bromhexine hydrochloride 8 mg

-Inactive ingredient(s):-
Magnesium stearate, lactose, Polyvinylpyrrolidone K-25 and Corn starch.

-MAL Number(s):-

MAL19911142XZ

Manufacturer:

HOVID Bhd.

Lot 56442, 7 1/2 Miles, Jalan Ipoh/Chemor, 31200 Chemor, Perak, Malaysia.

Product Registration Holder:

HOVID Bhd.

121, Jalan Tunku Abdul Rahman (Jalan Kuala Kangsar), 30010 Ipoh, Malaysia.

Date of revision:

8/3/2024

Serial Number:

NPRA(R4/1)070224/00016