

HOVID ADAMOL FILM-COATED TABLETS 500MG

Paracetamol Tablets (500mg)

What is in this leaflet

1. What Hovid Adamol Film-Coated Tablets is used for
2. How Hovid Adamol Film-Coated Tablets works
3. Before you use Hovid Adamol Film-Coated Tablets
4. How to use Hovid Adamol Film-Coated Tablets
5. While you are using it
6. Side effects
7. Storage and Disposal of Hovid Adamol Film-Coated Tablets
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of Revision

What Hovid Adamol Film-Coated Tablets is used for

Hovid Adamol Film-Coated Tablets is used for the relief of fever and the relief from mild to moderate pain including: headache, migraine, backache, musculoskeletal pain, period pain, pain of osteoarthritis, toothache, pain after dental procedures/tooth extraction, pain after vaccination and the discomfort from colds, influenza and sore throats.

How Hovid Adamol Film-Coated Tablets works

Hovid Adamol Film-Coated Tablets has analgesic (pain relief) and antipyretic (anti-fever) properties.

Before you use Hovid Adamol Film-Coated Tablets

- When you must not use it

Do not take Hovid Adamol Film-Coated Tablets if you are allergic to Paracetamol or Acetaminophen and other ingredients.

If you are not sure whether you should start taking this medicine, talk to your doctor or pharmacist.

- Before you start to use it

This product contains Paracetamol. Do not take other Paracetamol containing medicine at the same time.

Your doctor needs to know before you take Hovid Adamol Film-Coated Tablets if you have kidney or liver problem.

Hovid Adamol Film-Coated Tablets should also be taken with care when you take other medicines that affect the liver.

Discuss with your doctor or pharmacist first if you are pregnant or breastfeeding. Problems in human have not been reported when taking Hovid Adamol Film-Coated Tablets during pregnancy or breastfeeding.

- Taking other medicines

Tell your doctor or pharmacist before taking Hovid Adamol Film-Coated Tablets, if you are taking any other medicines, including any that you buy without a prescription from your pharmacy, supermarket or health food shop. The effects of some medicines may be affected if other medicines are used at the same time.

Inform your doctor if you are using Hovid Adamol Film-Coated Tablets in combination with the alcohol, liver enzyme inducers and hepatotoxic medications. Risk of liver toxicity increased with liver enzyme inducers and medicines that are known to be toxic to the liver.

How to use Hovid Adamol Film-Coated Tablets

Always take the medicine exactly as your doctor or pharmacist has told you. Read the direction on the labels carefully.

- How much to use

Adults (including the elderly) and children aged 12 years and over: 500mg to 1000mg paracetamol (1 to 2 tablets), taken every 4 to 6 hours as required.

Children under 12 years:

Not recommended for children under the age of 12 years.

Minimum dosing interval: 4 hours

Maximum daily dose: 4000mg (8 tablets)

- When to use it

Follow the direction label carefully and as instructed by your doctor or pharmacist.

- How long to use it

Follow the direction label carefully and as instructed by your doctor or pharmacist.

- If you forget to use it

Take the missed dose as soon as you remember. If it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to.

Do not take a double dose to make up for the dose that you missed. This may increase the chance of you getting unwanted side effects.

HOVID ADAMOL FILM-COATED TABLETS 500MG

Paracetamol Tablets (500mg)

- If you use too much (overdose)

Immediately call your doctor or go to the emergency department at your nearest hospital if you or anyone else has taken a large amount of this medication.

Symptoms of overdosage of Hovid Adamol Film-Coated Tablets can be seen in the first 24 hours. Among the symptoms are:

- pallor
- nausea
- vomiting
- anorexia
- abdominal pain

Liver damage (with a symptoms of pain, tenderness, and/ or swelling in the upper abdominal area) may become more apparent 12 to 48 hours after the intake of the medicine.

In severe poisoning, liver failure may progress to brain malfunction or damage, coma and death.

Prompt treatment is essential in the management of Hovid Adamol Film-Coated Tablets overdosage.

While you are using it

- Things you must do

Take the medication exactly as your doctor prescribed.

- Tell all the doctors, dentists and pharmacists treating you that you are taking Hovid Adamol Film-Coated Tablets.
- Tell your doctor immediately if you become pregnant while taking this medication.

- Things you must not do

Do not give Hovid Adamol Film-Coated Tablets to anyone else, even if their symptoms seem similar to yours.

Do not take your medicine to treat any other complaints unless your doctor or pharmacist tells you to.

- Things to be careful of

Avoid drinking alcohol while taking this medication.

Side effects

All medicines may have some unwanted side effects. Unwanted side effects do not always occur in every person. Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking Hovid Adamol Film-Coated Tablets.

- Allergy alert: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash.
- These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of Hovid Adamol Film-Coated Tablets

- Storage

Store below 30°C. Protect from light.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

Hovid Adamol Film-Coated Tablets is white to off white colored, circular, biconvex film coated tablets with break-line on one side and embossed with 'H' on other side.

- Ingredients

- Active ingredient(s)

Paracetamol

- Inactive ingredients

Pregelatinized Starch
Sodium Benzoate Povidone
K-30
Colloidal Anhydrous Silica Purified
Talc
Magnesium Stearate Wincoat
WT-TR-5002 Clear
Transparent Purified Water

- MAL number:

Manufacturer

RV Lifesciences Limited
Plot No. H-19, MIDC Waluj,
Chhatrapati Sambhajnagar, 431133,
India.

Product Registration Holder

Hovid Bhd.
121, Jalan Tunku Abdul Rahman
(Jalan Kuala Kangsar), 30010 Ipoh,
Perak, Malaysia

HOVID ADAMOL FILM-COATED TABLETS 500MG

Paracetamol Tablets (500mg)

Date of revision

13/04/2026

Serial Number

NPRA (R2) 20/094