




Batch No., Mfg. and Exp. dates
will be printed when used

For oral use.

Powder to be dissolved in water.

Directions for reconstitution: Loratyn Dry Syrup 1%w/w is administered by reconstituting the powder with 150 ml to 200 ml water. One sachet of Loratyn Dry Syrup 1% w/w provides 5 mg of Loratadine upon reconstitution and it should be taken immediately after preparation.

Refer package insert for dosage and other information.

Product Registration Holder: **HOVID Bhd.**
121, Jalan Tunku Abdul Rahman (Jalan Kuala Kangsar),
30010 Ipoh, Perak, Malaysia.

Manufactured by: **HOVID Bhd.**
Lot 56442, 7 1/2 Miles, Jalan Ipoh / Chemor,
31200 Chemor, Perak, Malaysia.

Controlled
Medicine

NON-DROWSY
LORATYN
DRY SYRUP 1% w/w

Loratadine Dry Syrup 1 %

- ✓ **ALLERGIC RHINITIS**
- ✓ **ALLERGIC DERMATOLOGIC DISORDER**
- ✓ **CHRONIC URTICARIA**

30 sachets x 0.5g

Raspberry Flavour

hovid

Each sachet contains:
Loratadine Micronised 1% w/w

Store below 30°C.
Keep in a dry place.

Keep out of reach of children.
Jauhi daripada kanak-kanak.

Reg. No.: MALxxxxxxAZ



VULORfa-PC3

Batch No.:
Mfg. Date :
Exp. Date :

NON-DROWSY
LORATYN
DRY SYRUP 1% w/w

Loratadine Dry Syrup 1 %

- ✓ **ALLERGIC RHINITIS**
- ✓ **ALLERGIC DERMATOLOGIC DISORDER**
- ✓ **CHRONIC URTICARIA**

30 sachets x 0.5g

Raspberry Flavour

hovid

hovid

- ✓ **ALLERGIC RHINITIS**
- ✓ **ALLERGIC DERMATOLOGIC DISORDER**
- ✓ **CHRONIC URTICARIA**

NON-DROWSY
LORATYN
DRY SYRUP 1% w/w
Loratadine Dry Syrup 1 %