
 e-Banner	Project: CO-0057288	Document: PPC-0026339	Version: 2
	Site Code: 62000000085275	Operator: MNB13223	Date/Time Created: 30.May.2023 15:28 GMT+1

Directions for making up the suspension: Invert and shake bottle to loosen powder. Add volume of water indicated in the package leaflet. Invert and shake well. Alternatively, fill the bottle with water to just below the mark on bottle label. Invert and shake well, then top up with water to the mark. Invert and shake again. Allow to stand for 5 minutes to ensure full dispersion. Shake well before taking each dose.

Controlled Medicine  
Manufactured by:  
Glaxo Wellcome Production  
Mayenne, France  
PRH in Malaysia:  
GlaxoSmithKline  
Pharmaceutical Sdn Bhd  
Petaling Jaya, Malaysia  
PRH in Brunei:  
Zuellig Pharma (B) Sdn Bhd  
BSB, Brunei Darussalam

BN  
MAN  
EXP




## Augmentin

amoxicillin trihydrate - potassium clavulanate

Twice daily dosing

**228 mg/5 ml**

70 ml suspension in powder form



When reconstituted, each 5 ml contains 200 mg amoxicillin (as amoxicillin trihydrate) and 28.5 mg clavulanic acid (as potassium clavulanate). Contains aspartame. Unsuitable for phenylketonurics.

To be taken as directed by the physician. Not to be used in patients with known hypersensitivity to penicillin.

**Keep out of the reach of children. Jauhkan daripada kanak-kanak.**

Store in a dry place at below 30°C in the original packaging to protect from moisture.

Once reconstituted, the suspension must be stored in a refrigerator (2°C to 8°C) and used within 7 days. Do not freeze.

For oral use only.  
MAL19886950ARZ BRU08030024P  
Trade marks are owned by or licensed to the GSK group of companies.  
©2023 GSK group of companies or its licensor.

62000000085275