

e-Banner	Project: CO-0057288	Document: PPC-0026341	Version: 2
	Site Code: 62000000085291	Operator: BT611974	Date/Time Created: 01.Jun.2023 05:21 GMT+1

BN
MAN
EXP



Directions for making up the suspension:
Invert and shake bottle to loosen powder.
Add volume of water indicated in the package leaflet. Invert and shake well.
Alternatively, fill the bottle with water to just below the mark on bottle label. Invert and shake well, then top up with water to the mark. Invert and shake again. Allow to stand for 5 minutes to ensure full dispersion. Shake well before taking each dose.
Controlled Medicine
Manufactured by:
Glaxo Wellcome Production
Meyenne, France
PRIH in Malaysia:
GlaxoSmithKline Pharmaceutical Sdn Bhd
Petaling Jaya, Malaysia
PRIH in Brunei:
Zuellig Pharma (B) Sdn Bhd
BSB, Brunei Darussalam



Augmentin

amoxicillin trihydrate - potassium clavulanate

Twice daily dosing

457 mg/5 ml

35 ml suspension in powder form



SUGAR FREE
Mixed Fruit Flavour

When reconstituted, each 5 ml contains 400 mg amoxicillin (as amoxicillin trihydrate) and 57 mg clavulanic acid (as potassium clavulanate). Contains aspartame. Unsuitable for phenylketonurics.
To be taken as directed by the physician.
Not to be used in patients with known hypersensitivity to penicillin.
Keep out of the reach of children.
Jauhkan daripada kanak-kanak.
Store in a dry place at below 30°C in the original packaging to protect from moisture. Once reconstituted, the suspension must be stored in a refrigerator (2°C to 8°C) and used within 7 days.
Do not freeze.
For oral use only.
MAL19986951ARZ BRU08030025P
Trade marks are owned by or licensed to the GSK group of companies.
©2023 GSK group of companies or its licensor.

62000000085291