

ARTWORK SIZE:
40 mm x 40.426 mm

IRON SUCROSE INJECTION USP 100 mg/ 5mL		5 mL
Referis		
Composition: Each 5mL contains: Iron(III) Hydroxide Sucrose Complex Equivalent to elemental iron... 100 mg Water for Injections Ph.Eur q.s. to 5 mL Sodium Hydroxide approximately... 7.5mg	For Intravenous Use Only Single use Ampoule CONTROLLED MEDICINE Reg. No. :MAL17085002AZ Mfg. Lic. No. :616	
Dosage & Administration: Refer Package insert.	Batch No. :	Unvarnished Area 13 x 10 mm
Storage: Store below 30°C. Do not freeze.	Mfg. Date :	
Keep medicine out of reach of children. Jauhkan daripada kanak-kanak	Exp. Date :	
Referis Injection is for intravenous injection. The injection must be diluted with 0.9% w/v Sodium chloride solution, to a concentration of 1 mg of elemental iron per mL. To be administered under physician's direct supervision.		
Precaution : Strict aseptic administration to be followed. Discard the unused solution. Caution : Before using check for absence of sediments. The injection should be discarded if any sediment appears.	Manufactured by: MAIVA PHARMA PRIVATE LIMITED No. 32, Sipcot Industrial Complex, Phase - 1, Hosur-635126, Tamil Nadu, INDIA.	
Osmolarity : Not less than 1150 and not more than 1350 mOsmol per liter.	Averroes Pharmaceuticals SDN BHD, 03-08-01 & 03-09-01, Block 3, Persint Alami, Worldwide Business center II, Persiaran Akuatik, Seksyen 13, 40100 Shah Alam, Selangor.	MY/LB/REF/02 PLC112/01/APD

Enlarge Size

IRON SUCROSE INJECTION USP 100 mg/ 5mL		5 mL
Referis		
Composition: Each 5mL contains: Iron(III) Hydroxide Sucrose Complex Equivalent to elemental iron... 100 mg Water for Injections Ph.Eur q.s. to 5 mL Sodium Hydroxide approximately... 7.5mg	For Intravenous Use Only Single use Ampoule CONTROLLED MEDICINE Reg. No. :MAL17085002AZ Mfg. Lic. No. :616	
Dosage & Administration: Refer Package insert.	Batch No. :	Unvarnished Area 13 x 10 mm
Storage: Store below 30°C. Do not freeze.	Mfg. Date :	
Keep medicine out of reach of children. Jauhkan daripada kanak-kanak	Exp. Date :	
Referis Injection is for intravenous injection. The injection must be diluted with 0.9% w/v Sodium chloride solution, to a concentration of 1 mg of elemental iron per mL. To be administered under physician's direct supervision.		
Precaution : Strict aseptic administration to be followed. Discard the unused solution. Caution : Before using check for absence of sediments. The Injection should be discarded if any sediment appears.	Manufactured by: MAIVA PHARMA PRIVATE LIMITED No. 32, Sipcot Industrial Complex, Phase - 1, Hosur-635126, Tamil Nadu, INDIA.	
Osmolarity : Not less than 1150 and not more than 1350 mOsmol per liter.	Averroes Pharmaceuticals SDN BHD, 03-08-01 & 03-09-01, Block 3, Persint Alami, Worldwide Business center II, Persiaran Akuatik, Seksyen 13, 40100 Shah Alam, Selangor.	MY/LB/REF/02 PLC112/01/APD