

Tebonin® forte Tablet 120mg

EGb 761® (Each tablet contains Patented Standardised Ginkgo biloba extract 120mg equivalent to dry leaf 6000mg)

Description & Composition

Tebonin® forte Tablet 120mg is presented as yellow and oval film-coated tablets in blister strip packing.

One film-coated tablet contains:

Active ingredient: EGb 761®, a dry extract from Ginkgo biloba leaves (50:1) 120mg. EGb 761® is standardised to 24% flavone glycosides, 6% terpene lactones & less than 5ppm ginkgolic acids.

Indications

Traditionally used to improve blood circulation and for general well-being.

General Warnings

As the use of Ginkgo may increase the tendency of bleeding, please consult your physician/pharmacist if you are on or intend to start using any other medicines and before you undergo any surgical/dental procedure. (Memandangkan Ginkgo boleh meningkatkan kemungkinan pendarahan, sila rujuk kepada doktor / ahli farmasi sekiranya anda sedang atau akan menggunakan ubat lain dan sebelum prosedur pembedahan / dental dijalankan).

It cannot be excluded that the application of Ginkgo preparations promotes the occurrence of seizures in epileptic patients.

Patients with rare hereditary galactose intolerance, lactase deficiency or glucose- galactose malabsorption should not take Tebonin® forte Tablet 120mg.

This is a Traditional Medicine.

Keep out of reach of children.

Jauhi daripada kanak-kanak.

Please consult your pharmacist/doctor before taking this product.

Contraindications

Tebonin® forte Tablet 120mg should not be used by patients with known hypersensitivity to Ginkgo extracts or to any of the excipients and is contraindicated in pregnancy and lactating women. Tebonin® forte Tablet 120mg is not recommended for use in adolescents aged from 12 to 18 years due to insufficient investigational data.

Adverse effects

In very rare cases, mild gastro-intestinal disorders, headache or allergic skin reactions (reddening, swelling, itching) have been observed following intake of Tebonin® forte Tablet 120mg. There have been very rare reports of bleeding, following the consumption of Ginkgo biloba. However, correlations between Tebonin® forte Tablet 120mg and bleeding have not been proven.

Drug Interactions

Specific clinical studies did not point to any interactions with acetylsalicyclic acid (aspirin) and warfarin. However, interaction with drugs inhibiting blood coagulation cannot be excluded.

Dosage and Administration

For adults, the normal dosage is 1 tablet of Tebonin® forte Tablet 120mg twice daily.

The intake of Tebonin® forte Tablet 120mg is independent of meals.

Expiry: 4 years from the date of manufacture

Pack size: 30's, 60's, 90's

Keep in a dry place below 30°C, protect from light and moisture

Date of revision: October 2023

Manufactured by



Dr. Willmar Schwabe
From Nature. For Health.

Dr. Willmar Schwabe GmbH & Co.KG
Willmar-Schwabe-Str. 4, D-76227 Karlsruhe, Germany



Product Registration Holder:

A. Menarini Singapore Pte Ltd (994247-T)

B-18-2, Level 18, The Ascent Paradigm,
No. 1 Jalan SS7/26A, Kelana Jaya,
47301 Petaling Jaya, Selangor

Product Registration: MAL07051012T

