

SKUDEXA (TRAMADOL HYDROCHLORIDE 75MG / DEXKETOPROFEN 25MG) FILM-COATED TABLETS

Tramadol Hydrochloride 75mg

Dexketoprofen 25mg

What is in this leaflet

1. What SKUDEXA is used for
2. How SKUDEXA works
3. Before you use SKUDEXA
4. How to use SKUDEXA
5. While you are using it
6. Side effects
7. Storage and Disposal of SKUDEXA
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

1. **What SKUDEXA is used for**
SKUDEXA is used for the symptomatic short term treatment of moderate to severe acute pain in adults.

2. How SKUDEXA works

SKUDEXA contains the active substances tramadol hydrochloride and dexketoprofen.

Tramadol hydrochloride is a pain killer belonging to a group of medicines called opioids that act on the central nervous system. It relieves pain by acting on specific nerve cells of the brain and spinal cord.

Dexketoprofen is a pain killer and it belongs to a group of medicines called non-steroidal anti-inflammatory drugs (NSAIDs).

3. Before you use SKUDEXA

- When you must not use it

- If you are allergic to dexketoprofen, to tramadol hydrochloride or to any of the other ingredients of this medicine
- If you are allergic to acetylsalicylic acid or to other NSAID
- If you have suffered attacks of asthma, acute allergic rhinitis (a short period of inflamed lining of the nose), nasal polyps (lumps

in the nose due to allergy), urticaria (skin rash), angioedema (swollen face, eyes, lips, or tongue, or respiratory distress) or wheezing in the chest after taking acetylsalicylic acid or other non-steroidal anti-inflammatory medicines

- If you have had photo allergic or phototoxic reactions (reddening and/or blistering of the skin exposed to sunlight) while taking ketoprofen (a NSAID) or fibrates (drugs used to lower the level of fats in the blood)
- If you have a peptic ulcer, stomach or bowel bleeding or if you have suffered in the past from stomach or bowel bleeding, ulceration or perforation, including that due to previous use of NSAIDs
- If you have chronic digestive problems (e.g. indigestion, heartburn)
- If you have bowel disease with chronic inflammation (Crohn's disease or ulcerative colitis)
- If you have serious heart failure, moderate or serious kidney problems or serious liver problems
- If you have a bleeding disorder or a blood clotting disorder
- If you are severely dehydrated (have lost a lot of body fluids) due to vomiting, diarrhoea or insufficient intake of fluids
- If you have acute poisoning with alcohol, sleeping pills, pain relievers, or medicines that affect mood and emotions
- If you are also taking monoamine oxidase inhibitors

(MAOIs) (certain medicines used for the treatment of depression) or have taken them in the last 14 days before treatment with this medicine

- If you have epilepsy or suffer from fits, because the risk of a fit may increase
- If you are breathing with difficulty
- If you are pregnant or breast feeding

- Before you start to use it

- If you have an allergy, or if you have had allergy problems in the past
- If you have kidney, liver or heart problems (hypertension and/or heart failure) as well as fluid retention, or have suffered from any of these problems in the past
- If you are taking diuretics
- If you have heart problems, previous stroke or think that you might be at risk of these conditions (for example if you have high blood pressure, diabetes or high cholesterol or are a smoker) you should discuss your treatment with your doctor; medicines such as this medicine may be associated with a small increased risk of heart attack (myocardial infarction) or stroke. Any risk is more likely with high doses and prolonged treatment. Do not exceed the recommended dose or duration of treatment
- If you are elderly: you may be more likely to suffer from side effects (see section 6). If any of these occur, consult your doctor immediately

SKUDEXA (TRAMADOL HYDROCHLORIDE 75MG / DEXKETOPROFEN 25MG) FILM-COATED TABLETS

Tramadol Hydrochloride 75mg

Dexketoprofen 25mg

- If you are a woman with fertility problems: this medicine may affect your fertility, therefore you should not take it if you are planning to become pregnant or you are having fertility tests
- Have a disorder in the formation of blood and blood cells
- If you have systemic lupus erythematosus or mixed connective tissue disease (immune system disorders that affect connective tissue)
- If you have suffered in the past from a chronic inflammatory disease of the bowel (ulcerative colitis, Crohn's disease)
- If you have or have suffered in the past from other stomach or bowel problems.
- If you have infection or chickenpox, since NSAIDs could worsen and mask the infection and its symptom, albeit rarely
- If you are taking other medicines that increase the risk of peptic ulcer or bleeding, e.g. oral steroids, some antidepressants (those of the SSRI type, i.e. Selective Serotonin Reuptake Inhibitors), drugs that
- prevent blood clots such as acetylsalicylic acid or anticoagulants such as warfarin. In such cases, consult your doctor before taking this medicine: he/she may want you to take an additional medicine to protect your stomach
- If you suffer from depression and you are taking antidepressant as some of them may interact with tramadol
- If you are taking other medicines containing the same active substances in this medicine, do not exceed the maximum daily doses of dexketoprofen or tramadol
 - If you think that you are addicted

to other pain relievers (opioids)

- If you have consciousness disorders (if you feel that you are going to faint)
- If you are in a state of shock (cold sweat may be a sign of this)
- If you have increased pressure in the brain (possibly after a head injury or brain disease)
- If you have difficulty in breathing
- If you have porphyria..

Tramadol may lead to physical and psychological addiction. When this medicine is taken for a long time, its effect may decrease, so that higher doses have to be taken (tolerance development). In patients with a tendency to abuse medicines or who are dependent on medicines, treatment with SKUDEXA should only be carried out for short periods and under strict medical supervision. Talk to your doctor if you experience any of the following symptoms while taking SKUDEXA: extreme fatigue, lack of appetite, severe abdominal pain, nausea, vomiting or low blood pressure. This may indicate that you have adrenal insufficiency (low cortisol levels). If you have these symptoms, contact your doctor, who will decide if you need to take hormone supplement.

Tramadol is transformed in the liver by an enzyme. Some people have a variation of this enzyme and this can affect people in different ways. In some people, they may not get enough pain relief but other people are more likely to get serious side effects. If you notice any of the following side effects, you must stop taking this medicine and seek immediate medical advice: slow or

shallow breathing, confusion, sleepiness, small pupils, feeling or being sick, constipation, lack of appetite.

There is a small risk that you may experience a so-called serotonin syndrome that can occur after having taken tramadol in combination with certain antidepressants or tramadol alone. Seek medical advice immediately if you have any of the symptoms related to this serious syndrome

Tell your doctor if any of these problems occurs during SKUDEXA treatment or if they applied in the past.

- Taking other medicines

Tell your doctor if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription. Some medicines should not be taken together and others may need their doses to be altered when taken together.

4. How to use SKUDEXA

-How much to use

Always use this medicine exactly as your doctor has told you. Check with your doctor if you are not sure.

The lowest effective dose should be used for the shortest duration necessary to relieve symptoms. If you have an infection, consult a doctor without delay if symptoms (such as fever and pain) persist or worsen.

The dose of SKUDEXA that you need depends on the type, severity and duration of your pain. Your doctor will tell you how many tablets you must take daily, and for how long.

The recommended dose is generally 1 film-coated tablet (corresponding to 75 mg of tramadol hydrochloride and 25 mg of dexketoprofen) every 8

SKUDEXA (TRAMADOL HYDROCHLORIDE 75MG / DEXKETOPROFEN 25MG) FILM-COATED TABLETS

Tramadol Hydrochloride 75mg
Dexketoprofen 25mg

hours, with no more than 3 film-coated tablets daily (corresponding to 225 mg of tramadol hydrochloride and 75 mg of dexketoprofen) and not exceeding 5 days of treatment.

Use in children and adolescents:

SKUDEXA is not suitable for children and adolescents.

Elderly patients:

If you are aged 75 years or over, your doctor may recommend prolonging the dosage interval because your body may handle the drug more slowly.

Severe liver or kidney disease (insufficiency)/dialysis patients:

Patients with severe liver and/or kidney insufficiency should not take SKUDEXA.

In case of renal dysfunction, if in your case the insufficiency is mild, your doctor may recommend prolonging the dosage interval.

In case of hepatic dysfunction, if in your case the insufficiency is mild or moderate, your doctor may recommend prolonging the dosage interval.

Swallow the tablet with a sufficient amount of fluid (preferably with a glass of water).

Food delays the absorption of SKUDEXA, so for a faster effect take the tablet at least 30 minutes before meals. The score line is to help you break the tablet if you have difficulty swallowing it whole.

- When to use it

Please refer to "How much to use" section.

- How long to use it

Please refer to "How much to use"

section.

- If you forget to use it

Do not take a double dose to make up for a forgotten tablet. Take the next regular dose when it is due (see section 4 "How to use SKUDEXA").

- If you use too much (overdose)

If you use too much of this medicine, tell your doctor immediately or go to the emergency department of your nearest hospital. Please remember to take this medicine pack or this leaflet with you.

The symptoms of an overdose of this medicine are:

- Vomiting, loss of appetite, stomach pain, drowsiness, dizziness/spinning sensation, disorientation, headache (for dexketoprofen)
- Contraction of the pupil, vomiting, heart failure, loss of consciousness, convulsions and difficulty in breathing (for tramadol). Serotonin syndrome has also been reported.

5. While you are using it

- Things you must do

Take your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are taking SKUDEXA

Tell your doctor immediately if you become pregnant while taking this medication

You must talk to a doctor if you do not feel better or if you feel worse.

- Things you must not do

Do not take any new medicines without consulting your doctor.

Do not give SKUDEXA to anyone

else, even if they have the same symptoms or condition as you.

- Things to be careful of

Use of SKUDEXA is not recommended with

- Acetylsalicylic acid, corticosteroids or other anti-inflammatory drugs;
- Warfarin, heparin or other medicines used to prevent blood clots;
- Lithium, used to treat certain mood disorders;
- Methotrexate, used for rheumatoid arthritis and cancer;
- Hydantoins and phenytoin, used for epilepsy;
- Sulfamethoxazole, used for bacterial infections; Monoamine oxidase inhibitors (MAOIs) (medicines for the treatment of depression).

Use of SKUDEXA requires precautions with:

- ACE inhibitors, diuretics, beta-blockers and angiotensin II antagonists, used for high blood pressure and heart problems
- Pentoxifylline, used to treat chronic venous ulcers
- Zidovudine, used to treat viral infections
- Chlorpropamide and glibenclamide used for diabetes
- Aminoglycoside antibiotics, used to treat bacterial infections.

Concomitant use of SKUDEXA and sedative medicines such as benzodiazepines or related drugs increases the risk of drowsiness, difficulties in breathing (respiratory depression), coma and may be life-threatening. Because of this, concomitant use should only be considered when other treatment options are not possible. However if

SKUDEXA (TRAMADOL HYDROCHLORIDE 75MG / DEXKETOPROFEN 25MG) FILM-COATED TABLETS

Tramadol Hydrochloride 75mg
Dexketoprofen 25mg

your doctor does prescribe SKUDEXA together with sedative medicines the dose and duration of concomitant treatment should be limited by your doctor. Please tell your doctor about all sedative medicines you are taking, and follow your doctor's dose recommendation closely. It could be helpful to inform friends or relatives to be aware of the signs and symptoms stated above. Contact your doctor when experiencing such symptoms.

Use with SKUDEXA requires care with:

- Quinolone antibiotics (e.g. ciprofloxacin, levofloxacin) used for bacterial infections;
- Ciclosporin or tacrolimus, used to treat immune system diseases and in organ transplant;
- Streptokinase and other thrombolytic or fibrinolytic medicines, i.e. medicines used to break up blood clots;
- Probenecid, used in gout;
- Digoxin, used to treat chronic heart failure;
- Mifepristone, used to terminate a pregnancy;
- Antidepressants of the selective serotonin reuptake inhibitors type (SSRIs);
- Anti-platelet agents used to reduce platelet aggregation and the formation of blood clots;
- Tenofovir, deferasirox, pemetrexed.

The pain-relieving effect of tramadol may be reduced, and the length of time it acts may be shortened, if you also take medicines containing: Carbamazepine (for epileptic fits); Buprenorphine, nalbuphine, or pentazocine (pain relievers); Ondansetron (prevents nausea).

The risk of side effects increases:

- if you take tranquilizers, sleeping pills, other pain relievers such as morphine and codeine (also as cough medicine), or alcohol while you are taking SKUDEXA. You might feel drowsier or feel that you might faint. If this happens tell your doctor
- if you are taking medicines which may cause convulsions (fits), such as certain antidepressants or antipsychotics. The risk of having a fit may increase if you take SKUDEXA at the same time. Your doctor will tell you whether SKUDEXA is suitable for you
- if you are taking certain antidepressants. SKUDEXA may interact with these medicines and you may experience Serotonin symptoms
- if you take anticoagulants (medicines for blood thinning), e.g. warfarin, together with this medicine. The effect of these medicines on blood clotting may be affected and bleeding may occur.

SKUDEXA with alcohol:

Do not drink alcohol during treatment with SKUDEXA as it may increase the effect of the medication.

Pregnancy, breast-feeding and fertility:

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine.

Dexketoprofen can cause kidney and heart problems in your unborn baby. It may affect your and your baby's tendency to bleed and cause labour to be later or longer than expected. From 20 weeks of pregnancy, dexketoprofen can cause kidney problems in your unborn

baby, that may lead to low levels of amniotic fluid that surrounds the baby (oligohydramnios) or narrowing of a blood vessel (ductus arteriosus) in the heart of the baby.

Tramadol is excreted into breast milk. The use of SKUDEXA is contraindicated in pregnancy as well as during breast-feeding.

Driving and using machines:

SKUDEXA may affect your ability to drive and handle machines, due to the possibility of dizziness, blurred vision or drowsiness as side effects of treatment. This applies particularly when SKUDEXA is taken with medicines that affect mood and emotions, or alcohol.

If you are affected, do not drive or use machines until the symptoms wear off.

This medicinal product contains less than 1 mmol sodium (23mg) per dosage unit, that is to say essentially "sodium-free".

6. Side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Very common side effects (may affect more than 1 in 10 people):

- nausea/feeling sick
- dizziness.

Common side effects (may affect up to 1 in 10 people):

- vomiting
- stomach pain
- diarrhoea
- digestive problems
- headaches
- drowsiness, fatigue

SKUDEXA (TRAMADOL HYDROCHLORIDE 75MG / DEXKETOPROFEN 25MG) FILM-COATED TABLETS

Tramadol Hydrochloride 75mg

Dexketoprofen 25mg

- constipation
- dry mouth
- increased sweating.

Uncommon side effects (may affect up to 1 in 100 people):

- increase in the number of blood platelets
- effects on the heart and blood circulation (pounding of the heart, fast heartbeat, feeling faint or collapse), low blood pressure. These adverse effects may occur particularly when patients are in an upright position or under physical strain.
- High or very high blood pressure
- swelling in the voice-box (laryngeal oedema)
- reduced potassium in the blood
- psychotic disorder
- swelling next to the eye
- shallow or slow breathing
- discomfort, abnormal feeling,
- blood in urine
- spinning sensation
- sleeplessness or difficulty falling asleep
- nervousness/anxiety
- flushing
- flatulence, tiredness
- pain
- feeling feverish and shivering, generally feeling unwell
- abnormal blood tests
- urge to vomit (retching)
- feeling of pressure in the stomach, bloating
- inflammation of the stomach, skin reactions (e.g. itching, rash)
- facial swelling

Rare side effects (may affect up to 1 in 1,000 people):

- swelling of the lips and throat
- peptic ulcer, peptic ulcer perforation or bleeding, which may be seen as vomiting blood or black stools
- prostate problems
- liver inflammation (hepatitis), liver damage
- acute kidney failure
- slow heartbeat
- epileptic fits
- allergic/anaphylactic reactions (e.g. difficulty breathing, wheezing, swelling of the skin) and shock (sudden circulatory failure)
- transient loss of consciousness (syncope)
- hallucinations
- water retention or swollen ankles,
- loss of appetite, changes in appetite
- acne
- back pain
- passing urine frequently, or less than normal, with difficulty or pain
- menstrual disorders
- abnormal sensations (e.g. itching, tingling, numbness)
- trembling, muscle twitches, uncoordinated movement, weak muscles,
- confusion
- sleep disorders and nightmares
- disturbed perception
- blurred vision, contraction of the pupil
- shortness of breath.
- Psychological side effects may occur after treatment with SKUDEXA. Their intensity and nature may vary (according to the patient's personality and length of

therapy):

- change in mood (mostly high spirits, occasionally irritation)
- changes in activity (slowing down but sometimes an increase in activity)
- being less aware
- less able to make decisions, which may lead to errors in judgement.

Worsening of asthma has been reported. When treatment is stopped abruptly signs of withdrawal may appear. Epileptic fits have occurred mainly at high doses of tramadol or when tramadol was taken at the same time as other medicines which may induce fits.

Very rare side effects (may affect up to 1 in 10,000 people):

- inflammation of the pancreas
- kidney problems
- reduced white blood cell count (neutropenia)
- fewer platelets in the blood (thrombocytopenia)
- open sores on skin, mouth, eyes and genital areas (Stevens Johnson and Lyell's syndromes)
- breathlessness due to narrowing of the airways
- ringing in the ears (tinnitus)
- sensitive skin
- sensitivity to light.

SKUDEXA (TRAMADOL HYDROCHLORIDE 75MG / DEXKETOPROFEN 25MG) FILM-COATED TABLETS

Tramadol Hydrochloride 75mg

Dexketoprofen 25mg

Not known (frequency cannot be estimated from the available data):

- speech disorders
- extreme pupil dilatation
- decrease in blood sugar levels.
- Serotonin syndrome, that can manifest as mental status changes (e.g. agitation, hallucinations, coma), and other effects, such as fever, increase in heart rate, unstable blood pressure, involuntary twitching, muscular rigidity, lack of
- coordination and/or gastrointestinal symptoms (e.g. nausea, vomiting, diarrhoea)
- Hiccup

Tell your doctor immediately if you experience symptoms of an allergic reaction such as swollen face, tongue and/or throat, and/or difficulty swallowing or hives together with difficulties in breathing, experience stomach/bowel side effects at the start of treatment (e.g. stomach pain, heartburn or bleeding), if you have previously suffered from any such side effects due to long-term use of anti-inflammatory drugs, and especially if you are elderly.

The most common side effects during treatment with SKUDEXA are nausea and dizziness, which occur in more than 1 out of 10 patients.

During treatment with NSAIDs, fluid retention and swelling (especially in the ankles and legs), increased blood pressure and heart failure have been reported.

Medicines such as SKUDEXA may be associated with a small increased risk of heart attack or stroke.

In patients with immune system disorders that affect connective tissue (systemic lupus erythematosus or mixed connective tissue disease), anti-inflammatory medicines may rarely cause fever, headache and neck stiffness.

You should see a doctor immediately if you experience symptoms of an allergic reaction such as swollen face, tongue and/or throat, and/or difficulty swallowing or hives together with difficulties in breathing.

Stop using SKUDEXA as soon as you notice the appearance of a skin rash, or any lesion inside the mouth or on mucous membranes, or any sign of an allergy.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.moh.gov.my (Consumer → Reporting Side Effects to Medicines (ConSERF or Vaccines (AEFI)

7. Storage and Disposal of SKUDEXA

- Storage

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and on the blister after EXP. The expiry date refers to the last day of that month.

Store below 30°C in the original package in order to protect from light.

- Disposal

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These

measures will help protect the environment.

8. Product Description

- What it looks like

Almost white to slightly yellow, oblong, film-coated tablets with a break-mark on one side and debossed "M" on the other side in plastic/aluminium push-through pill packs.

- Ingredients

- Active ingredient(s):

75 mg of tramadol hydrochloride and 25 mg dexketoprofen.

- Inactive ingredients:

Tablet core: microcrystalline cellulose, pregelatinised maize starch, croscarmellose sodium, sodium stearyl fumarate, anhydrous silica colloidal.

Film-coating: polyvinyl alcohol, titanium dioxide, Macrogol/PEG 3350, talc.

- MAL number(s):

MAL20126002AZ

9. Manufacturer

Menarini - Von Heyden GmbH
Leipziger Strasse 7-13
01097 Dresden
Germany

Product Registration Holder

A. Menarini Singapore Pte Ltd
B-18-2, Level 18, The Ascent
Paradigm, No 1, Jalan SS 7/26A,
Kelana Jaya, 47301 Petaling Jaya,
Selangor, Malaysia

10. Date of revision

14 April 2023

Serial Number

NPRA (R1/1) 16112023/199