

Front

255 mm

76 mm

ABOUT YOUR TREATMENT

You have been prescribed Lixiana®, an anticoagulant medicine, which thins your blood and helps prevent you from suffering blood clots. It is important that you take your medication exactly as instructed by your doctor.

- If you miss a dose, take it immediately and then continue the following day as normal – do not take double the prescribed dose in a single day.

- Do not start any other medications (including over the counter) without consulting your doctor.
- Do not stop taking Lixiana® without consulting your doctor as this can increase your risk of developing a blood clot.

WHEN TO SEEK MEDICAL ADVICE

RISK OF BLEEDING

Taking an anticoagulant medicine such as Lixiana® can increase your risk of bleeding. It is therefore important to be aware of the possible signs and symptoms of bleeding and to speak to your doctor **immediately** if you experience any of the following:

- Bruising or bleeding under the skin
- Blood in the urine
- Coughing up blood

- Vomiting blood or material that looks like ground coffee
- Nose bleeds or cuts that take a long time to stop bleeding
- Tar-coloured stools
- Dizziness or sudden headache
- Unexplained tiredness
- Abnormal vaginal bleeding, including heavier or prolonged menses

Please talk to your doctor if you experience any unusual symptoms.

PATIENT ALERT CARD

Lixiana®
film-coated tablets
edoxaban

Please keep this card with you at all times.

Present it to your healthcare professional, pharmacist, surgeon or dentist before any medical treatment or intervention.

179400001 P1/20 E1/2

2018MEN-0336-1

16

51 mm

51 mm

51 mm

51 mm

51 mm

Back

255 mm

76 mm

PATIENT INFORMATION

Patient name:

Date of birth:

In case of emergency, please contact:

Name:

Phone no:

TREATMENT INFORMATION
(To be completed by physician)

Lixiana® has been prescribed at a once-daily dose of: mg

Started on: / (mm/yy)

Blood type:

Other medications/conditions:

.....

.....

.....

PRESCRIBER INFORMATION

For more information or in case of emergency, please contact:

Physician's name:

Phone number, practice stamp:

Signature of physician:

INFORMATION FOR HEALTHCARE PROFESSIONALS

- Lixiana® is an oral anticoagulant factor Xa inhibitor.
- When an invasive procedure is required, Lixiana® should be stopped at least 24 hours beforehand, and appropriate caution exercised.
- Lixiana® may increase the risk of bleeding. In case of clinically significant bleeding, stop treatment immediately.

- Coagulation tests such as INR, prothrombin time (PT), or activated partial thromboplastin time (aPTT) are not a useful measure of the effect of Lixiana®. However, a calibrated anti-Factor Xa assay may help inform clinical decisions.

Please consult the package insert for more information.

 Daiichi-Sankyo

This document has been approved by HSA on 18-12-2020.

179400001 P1/20 E2/2

51 mm

51 mm

51 mm

51 mm

51 mm

Pantone Black C

Pantone 233 C

IPM no: 2018MEN-0336-1
Product name: Lixiana Patient Alert Card
Countries: MY SG
Dimension: 255 X 76 (Endformat 51 X 76) mm.
Version: 10
Date: 22 December 2020