


Front

255 mm					
76 mm	<p>ABOUT YOUR TREATMENT</p> <p>You have been prescribed Lixiana®, an anticoagulant medicine, which thins your blood and helps prevent you from suffering blood clots. It is important that you take your medication exactly as instructed by your doctor.</p> <ul style="list-style-type: none"> If you miss a dose, take it immediately and then continue the following day as normal – do not take double the prescribed dose in a single day. 	<ul style="list-style-type: none"> Do not start any other medications (including over the counter) without consulting your doctor. Do not stop taking Lixiana® without consulting your doctor as this can increase your risk of developing a blood clot. 	<p>WHEN TO SEEK MEDICAL ADVICE</p> <p>RISK OF BLEEDING</p> <p>Taking an anticoagulant medicine such as Lixiana® can increase your risk of bleeding. It is therefore important to be aware of the possible signs and symptoms of bleeding and to speak to your doctor immediately if you experience any of the following:</p> <ul style="list-style-type: none"> Bruising or bleeding under the skin Blood in the urine Coughing up blood 	<ul style="list-style-type: none"> Vomiting blood or material that looks like ground coffee Nose bleeds or cuts that take a long time to stop bleeding Tar-coloured stools Dizziness or sudden headache Unexplained tiredness Abnormal vaginal bleeding, including heavier or prolonged menses <p>Please talk to your doctor if you experience any unusual symptoms.</p>	<p style="text-align: center;">PATIENT ALERT CARD</p> <p style="text-align: center;">Lixiana® film-coated tablets edoxaban</p> <p style="text-align: center;">Please keep this card with you at all times.</p> <p style="text-align: center;">Present it to your healthcare professional, pharmacist, surgeon or dentist before any medical treatment or intervention.</p> <p style="text-align: right; font-size: small;">179400001 P1/20 E1/2</p>
	51 mm	51 mm	51 mm	51 mm	51 mm

Back

255 mm					
76 mm	<p>PATIENT INFORMATION</p> <p>Patient name:</p> <p>Date of birth:</p> <p>In case of emergency, please contact:</p> <p>Name:</p> <p>Phone no:</p>	<p>TREATMENT INFORMATION (To be completed by physician)</p> <p>Lixiana® has been prescribed at a once-daily dose of: mg</p> <p>Started on: / (mm/yy)</p> <p>Blood type:</p> <p>Other medications/conditions:</p>	<p>PRESCRIBER INFORMATION</p> <p>For more information or in case of emergency, please contact:</p> <p>Physician's name:</p> <p>Phone number, practice stamp:</p> <p>Signature of physician:</p>	<p>INFORMATION FOR HEALTHCARE PROFESSIONALS</p> <ul style="list-style-type: none"> Lixiana® is an oral anticoagulant factor Xa inhibitor. When an invasive procedure is required, Lixiana® should be stopped at least 24 hours beforehand, and appropriate caution exercised. Lixiana® may increase the risk of bleeding. In case of clinically significant bleeding, stop treatment immediately. 	<ul style="list-style-type: none"> Coagulation tests such as INR, prothrombin time (PT), or activated partial thromboplastin time (aPTT) are not a useful measure of the effect of Lixiana®. However, a calibrated anti-Factor Xa assay may help inform clinical decisions. <p>Please consult the package insert for more information.</p> <p style="text-align: center;"> Daiichi-Sankyo</p> <p>This document has been approved by HSA on dd-mm-yyyy.</p> <p style="text-align: right; font-size: small;">179400001 P1/20 E2/2</p>
	51 mm	51 mm	51 mm	51 mm	51 mm

Pantone Black C
 Pantone 233 C
 IPM no: 2018MEN-0336-1
 Product name: Lixiana Patient Alert Card
 Countries: MY SG
 Dimension: 255 X 76 (Endformat 51 X 76) mm.
 Version: 8
 Date: 28 September 2020