FINJUVE FOR MEN 2.275 MG/ML CUTANEOUS SPRAY, SOLUTION

Finasteride (2.275g/mL)

What is in this leaflet

1. What is FINJUVE used for
2. How FINJUVE works
3. Before you use FINJUVE
4. How to use FINJVUE
5. While you are using it
6. Side effects
7. Storage and Disposal of FINJUVE
8. Product Description
9. Manufacturer
10. Date of revision

What FINJUVE is used for

Finjuve contains the active substance finasteride. It is administered to the skin of the bald scalp using a spray applicator consisting of a bottle with a pump and cone. Finjuve is used for the treatment of mild to moderate male pattern hair loss (also known as androgenetic alopecia). Finjuve increases hair growth on the scalp and prevents further hair loss in men. Finjuve is only for use by men from 18 to 41 years of age

How FINJUVE works

Male pattern hair loss is a common condition thought to be caused by a combination of genetic factors and a particular hormone, dihydrotestosterone (DHT). In male pattern hair loss, the balding scalp contains increased amounts of DHT. This hormone is thought to contribute to shortening the length of time that the hair grows and to thinning of the hair. As the hair follicles become smaller (miniaturised), balding becomes apparent. Finjuve lowers the level of DHT in the scalp. This helps to reverse the balding process, leading to an increased hair growth and prevention of further hair loss

Before you use FINJUVE

- When you must not use it

- If you are allergic to finasteride or any of the other ingredients of this medicine.
- If you are a woman and are or may become pregnant.

- Before you start to use it

Talk to your doctor or pharmacist before using Finjuve

Possible transfer of Finjuve

This medicine may cause a male baby to be born with abnormalities of the sex organs if the active ingredient finasteride is absorbed through the skin of a pregnant woman. Avoid any contact between the treated area and a woman who is or may become pregnant. You should also tell her to avoid touching surfaces exposed to Finjuve. If contact with Finjuve occurs, the woman concerned should wash the affected body part quickly and thoroughly.

Children and adolescents must not come into contact with Finjuve. If contact with Finjuve occurs, the child or adolescent concerned should wash the affected body part quickly and thoroughly.

Effects on Prostate-Specific Antigen (PSA)

If you have a prostate-specific antigen (PSA) blood test done for prostate cancer screening, tell your doctor that you are using Finjuve as it may affect interpretation of the results.

Effect on male hormone dihydrotestosterone (DHT)

Finjuve makes the concentration of the male hormone DHT in blood go down and commonly even to below normal values. However, this occurs less frequently and also the decrease is lower than with finasteride tablets. Side effects of a sexual nature that are known for finasteride tablets may also occur with Finjuve but are less likely (see section 4). Therefore, adhere to the dose that your doctor has prescribed for you.

Breast cancer

Although breast cancer has not occurred in men treated with Finjuve in clinical studies, it has been reported during treatment with finasteride tablets. If you experience any changes in your breast tissue such as lumps, pain, enlargement or nipple discharge, contact your doctor as soon as possible.

Mood alteration and depression

Although mood alteration has not been observed in patients treated with Finjuve in clinical studies, it has been reported during treatment with finasteride tablets. If you experience symptoms such as low mood, depression or suicidal thoughts, contact your doctor for advice as soon as possible.

Children and adolescents

Finjuve should not be used by children or adolescents. There are no data demonstrating efficacy or safety of finasteride for children under the age of 18 years.

Pregnancy

Finjuve must not be used by women. Women who are or may become pregnant must avoid contact with the treated scalp or surfaces exposed to Finjuve. See section "Possible transfer of Finjuve" above. If direct contact with this medicine occurs, the woman should wash the affected body part quickly and thoroughly and ask her doctor for advice.

Driving and using machines

Finjuve has no influence on the ability to drive and use machines.

Finjuve contains ethanol

This medicine contains 25 mg ethanol (96 per cent) in each spray which is equivalent to 0.5 mg/microlitre (55 per cent). It may cause burning sensation on damaged skin.

- Taking other medicines

Tell your doctor or pharmacist if you are taking, have recently taken, or might take any other medicines.

Do not apply Finjuve if you are using other topical products, such as cosmetics, sunscreens or other medicines, in the same area.


How to use FINJUVE

- How much to use

Depending on how much of your scalp is bald, your doctor will prescribe between 1 and 4 sprays daily to be used on separate areas. Do not use more than 4 sprays daily. Finjuve is for cutaneous use. It is only to be used on the scalp.

Ensure hair and scalp are fully dry prior to application. Apply Finjuve on the scalp only by yourself. If more than one spray is prescribed, apply these to areas that do not overlap. Do not apply the solution to areas of the body other than the scalp. Once applied, leave Finjuve in place for at least 6 hours.


Components and assembly of the spray applicator



Assembly of the spray applicator



A. Align and press:
Align the cone with the pump button and firmly press into place



B. Correct assembly:
The spray applicator is correctly assembled when you hear a click after pressing the spray nozzle is located in the centre position of the cone



C Correct assembly
and the bottom of the pump button is lined up with the bottom of the cone without any gap



D Incorrect, gap.
Realign and press again. If you do not hear a click during assembly or you can see a gap between the bottom of the pump button and the cone, realign the components and press into place again.

Priming the pump

- After assembly of the spray applicator the pump must be primed for the first use. If

FINJUVE FOR MEN 2.275 MG/ML CUTANEOUS SPRAY, SOLUTION

Finasteride (2.275g/mL)

the spray applicator has not been used for 2 weeks or longer, the pump will need priming again. There is no need to reprime at each use.

- To prime the pump for the first time, fully press the pump four times with your thumb or index finger, directing the sprayed solution into a sink. Then rinse the sink with water. To reprime the pump after non-use for 2 weeks or longer fully press the pump one time only.
- Do not spray Finjuve towards your face.
- If solution is released during assembly or priming, clean surfaces where it may have been deposited.

Priming




Do no spray towards the face



Dose application

- There is no need to shake the bottle before use.
- Hold the spray applicator with the cone flat against the scalp to avoid dispersion of the solution in the air.
- Fully press the pump once for one spray.
- Move the cone to different areas of your scalp to apply any further doses according to the number of sprays prescribed by your doctor. Do not overlap and treat areas that have already been sprayed.
- After use, do not remove the cone from the pump. Put the spray applicator back in the box.
- Once applied, do not wash Finjuve off for at least 6 hours.

Cone flat against scalp



Move cone to the next area which does not overlap



Ensure that Finjuve does not come into contact with your hands or any other parts of the body. Wash immediately and thoroughly any part of the body exposed other than your scalp. If the cone becomes dirty, wipe it with a clean dry tissue. Safely throw away the used tissue and wash your hands thoroughly.

- When to use it

1-4 times a day as instructed by your doctor

- How long to use it

The bottle contains up to 180 sprays. The number of days of treatment depends on the prescribed dose, 1 to 4 sprays per day.
On the date you start treatment with Finjuve make a note in your calendar of the prescribed dose (1 to 4 sprays) and calculate when a new bottle is needed. Contact your doctor before your supply runs out so there is no break in your treatment. Do not use the bottle beyond 180sprays, because any remaining solution in the bottle may not give a full dose, which could limit the effect of your treatment.

Sprays per day	Days of treatment
1	180
2	90
3	60
4	45

- If you forget to use it)

If you forget to apply Finjuve, do not apply a double dose to make up for the forgotten application. Continue to use at the dose recommended by your doctor.

- If you use to much (overdose)

If you apply more Finjuve than recommended, talk to your doctor. Finjuve will not work faster or better if you apply it more than once a day, but side effects may be more likely.

While you are using it

- Things you must do

It may take 3 months for the treatment effect to develop. It is important to keep using Finjuve for as long as your doctor tells you. If you stop applying Finjuve, you are likely to lose the hair you have gained.

- Things you must not do

Do not use more than 4 sprays daily.

- Things to be careful of

Finjuve can be transferred through contact with fabrics, hands or other surfaces and objects. Avoid contact between the treated scalp and pillows, helmets, hats etc until the solution has dried. Finjuve can transfer from your body to others if they touch your treated scalp or other exposed surfaces. If contact with Finjuve occurs, the person should wash the affected body part quickly and thoroughly. Advise family members or others with access to the storage location of the contact precautions.

Side Effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.
Common (up to 1 in 10 people)
- Itching or redness of the scalp

Very common (> 1 in 10 people)
- Decrease of a male hormones (dihydrotestosterone) in blood

For other side effects, please refer to the package insert.

You may report side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers→ Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of FINJUVE

- Storage

Keep this medicine out of the sight and reach of children. Do not use this medicine after the expiry date which is stated on the box and the bottle label after EXP. Do not store above 30ºC. Finjuve contains alcohol and is therefore flammable. Avoid spraying the medicine near open flames or while smoking. Do not use Finjuve for longer than 6 months after first opening the bottle.

- Disposal

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use.

Product Description

- What is looks like

Each spray delivers 50 microlitres, which contain 114 micrograms of finasteride. It is a colourless, clear and slightly viscous solution.

- Ingredients

- Active ingredients(s)
2.275 mg of finasteride per ml
-Inactive ingredients
ethanol (96 per cent), purified water, propylene glycol, hydroxypropyl chitosan

-MAL number

MAL25046013AZ

Manufacturer

Almirall Hermal GmbH
Scholtzstraβe 1 and 3
21465 Reinbek
Germany

Product Registration Holder

A. Menarini Singapore Pte Ltd
B-18-2, Level 18, The Ascent Paradigm,
No 1, Jalan SS 7/26A, Kelana Jaya, 47301
Petaling Jaya, Selangor, Malaysia

Date of revision

15/04/2026

Serial Number

NPRA (R1) 24/040