

TARLIGE® FILM-COATED TABLET

Mirogabalin besilate (2.5mg, 5mg, 10mg & 15mg)

What is in this leaflet

1. What is TARLIGE® used for
2. How TARLIGE® works
3. Before you use TARLIGE®
4. How to use TARLIGE®
5. While you are using it
6. Side effects
7. Storage and Disposal of TARLIGE®
8. Product Description
9. Manufacturer
10. Date of revision

What TARLIGE® is used for

It is prescribed to people with neuropathic pain.

How TARLIGE® works

This medicine belongs to a group called neuropathic pain therapeutic agents. This medicine suppresses neurotransmitter release and exerts analgesic effect by suppressing influx of calcium ion in nerves.

Before you use TARLIGE®

- When you must not use it

The following persons cannot take this medicine:

- Persons with a history of hypersensitivity to any of the active ingredients and excipients in Tarlige

- Before you start to use it

The following patients may need to be careful when using this medicine: Tell your doctor or pharmacist before you start this medicine

- Persons with renal dysfunction
- Pregnant or possibly pregnant women
- Breastfeeding women

- Taking other medicines

Some drugs should be taken with care of this medicine. If you are taking any other medicine or new use of this medicine, consult with your doctor or pharmacist

How to use TARLIGE®

- How much to use

The dose and number of doses will be determined by your doctor according to your symptoms.

- For elderly people and people with kidney problems, the amount and number of doses may be adjusted.

- Usually, the amount and number of doses for adults are as follows. The effective dose may be adjusted according to your age and symptoms.

	Initial dose	Effective dose
--	--------------	----------------

One dose	5mg	10 – 15mg
Number of doses	Twice a day	

After beginning administration, the dose will be increased by 5mg with an interval of 1 week or longer

- When to use it

Twice a day

- How to take this medicine?

Take with glass of water or lukewarm water.

- How long to use it

It will be determined by your doctor. This medicine may cause insomnia, nausea, diarrhea, loss of appetite, etc. if you stop taking this medicine or decrease the dosage at your own discretion. It is important to continue taking this medicine as instructed.

- If you forget to use it

You should never take two doses at one time.

If you miss a dose, take the missed dose as soon as possible. If it is almost time for the next dose, skip the missed dose and follow your regular dosing schedule.

- If you use too much (overdose)

This medicine may cause symptoms such as euphoric mood, dysarthria, headache, dysphagia (difficulty swallowing food and water), sarcopenia, arthritis (swelling of joints), joint swelling and lack of strength. If any of these symptoms occur at the same time, consult a doctor immediately

While you are using it

- Things you must do

- Pay attention to the risk of falling and fracture due to these symptoms, especially in elderly people.
- Pay attention to obesity because it may increase body weight. Weight gain is common, especially if the dose is increased or if you are taking this medicine for a long time. Measure body weight regularly. If you start to gain weight, consult with your doctor. Improvements in diet and exercise may be recommended.

- Things you must not do

- This medicine may cause dizziness, somnolence or loss of consciousness. Avoid operating dangerous machinery, such as driving a car.
- Sudden discontinuation of this

medicine may cause symptoms such as insomnia, nausea, diarrhea and loss of appetite. If you stop taking this medicine, the dosage should be gradually decreased over time. Do not reduce the dose or stop taking this medicine at your own discretion.

- Alcoholic beverages may affect this medicine. Refrain from drinking alcoholic beverages.

- Things to be careful of

- This medicine may cause symptoms such as lazy eye, abnormal vision, blurred vision, double vision. If any of these symptoms occur, consult with your doctor.
- If you are pregnant or possibly pregnant, consult with your doctor.
- If you are breastfeeding, consult with your doctor.
- If you visit other doctors or purchase other medicines at pharmacies, etc., inform your doctor or pharmacist that you are taking this medicine.

Side Effects

This section describes the major adverse reactions and the major subjective symptoms. In the case of adverse reactions, it is common for some of the main subjective symptoms described for each significant adverse reaction to occur at similar times.

If any of these symptoms occur, consult with your doctor or pharmacist immediately.

Clinically significant adverse reactions	Main subjective symptoms
Dizziness	Floating feeling, lightheadedness, and spinning sensation
Somnolence	Drowsiness, sleepiness
Loss of consciousness	Losing consciousness
Hepatic dysfunction	Fatigability, feeling listlessness, weakness, nausea, decreased appetite
Renal impairment	Decreased urine output, edema, feeling listlessness

The above subjective symptoms are arranged according to the site of the adverse reaction as follows. If you notice these symptoms, see the Table for Significant Adverse Reactions

Site	Subjective symptoms
Whole body	Floating feeling, lightheadedness, fatigability, general

TARLIGE® FILM-COATED TABLET

Mirogabalin besilate (2.5mg, 5mg, 10mg & 15mg)

	malaise, weakness
Head	Spinning sensation, drowsiness, sleepiness, loss of consciousness
Mouth and throat	Nausea
Urine	Decreased urine output

Manufacturer

Takatsuki Plant, TAIYO Pharma Tech Co., Ltd,
4-38 Aketacho, Takatsuki, Osaka, Japan

Product Registration Holder

A. Menarini Singapore Pte Ltd
B-18-2, Level 18, The Ascent Paradigm,
No 1, Jalan SS 7/26A, Kelana Jaya, 47301
Petaling Jaya, Selangor, Malaysia

Date of revision

23/09/2025

Serial Number

(R1) 25/024

You may report side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers→ Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of TARLIGE®**- Storage**

- Store at below 30°C away from direct sunlight and moisture.
- Store the solution out of reach of children.

- Disposal

- Definitely don't give it to anyone else.
- Consult with your pharmacy or medical institution on how to dispose of unused medicines

Product Description**- What it looks like**

Tarlige Tablets 2.5mg: Peach round-shaped film-coated tablets

Tarlige Tablets 5mg: Grayish-red oblong-shaped filmcoated tablets with debossed line for scoring

Tarlige Tablets 10mg: Peach oblong-shaped film-coated tablets with debossed line for scoring

Tarlige Tablets 15mg: Grayish-red oblong-shaped filmcoated tablets with debossed line for scoring

- Ingredients

- Active ingredients(s)

Mirogabalin besilate

-Inactive ingredients

D-mannitol, microcrystalline cellulose, carmellose calcium, tocopherol, citric acid hydrate, magnesium aluminometasilicate, magnesium stearate, hypromellose, titanium oxide, talc, yellow ferric oxide, red ferric oxide

-MAL number

MALXXXXXXXXA