

DYNA YELLOW LOTION

Acriflavine (0.2% w/v)

What is in this leaflet

1. What Dyna Yellow Lotion is used for
2. How Dyna Yellow Lotion works
3. Before you used Dyna Yellow Lotion
4. How to use Dyna Yellow Lotion
5. While you are using it
6. Side effects
7. Storage and disposal of Dyna Yellow Lotion
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Dyna Yellow Lotion is used for

Indicated as an antiseptic for minor cuts, wounds and burns.

How Dyna Yellow Lotion works

The acriflavine hydrochloride is a disinfectant Capable of inhibiting the growth or reproduction of bacteria and have antifungal activity.

Before you use Dyna Yellow Lotion

- When you must not use it

Pregnancy and lactation

Ask your doctor or pharmacists for advice before using Dyna Yellow Lotion.

- Before you start to use it

Tell your doctor if you are hypersensitive to Acriflavine or other ingredients.

- Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

How to use Dyna Yellow Lotion

- How much to use

Apply to the affected part as required.

The directions given may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

- When to use it

Use as directed by your doctor or pharmacist.

- How long to use it

Continue taking Dyna Yellow Lotion for as long as your doctor recommends.

- If you forget to use it

Take the missed dose as soon as you remember. If it almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

The preparation is well tolerated following the recommended directions.

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using it

- Things you must do

Use your medicine exactly as your doctor has told you.

Tell the doctor, dentist and pharmacist treating you that you are taking Dyna Yellow Lotion.

- Things you must not do

Do not take any new medicines without consulting your doctor or pharmacist.

Do not give Dyna Yellow Lotion to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

Avoid getting into eyes or on mucous membranes.

Discontinue use if excessive irritation of the skin develops.

Side effects

With the recommended dosage and directions, side effects are unlikely to occur.

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine. You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website nptra.gov.my (Consumers → Reporting

Side Effects to Medicines (ConSERF) or Vaccines (AEFI).

Storage and Disposal of Dyna Yellow Lotion

- Storage

Keep out of reach of children to avoid accidental poisoning.

Protect from light

Store in dry place below 30°C

Keep the container tightly closed

For external use only

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

Yellow colour lotion with natural fragrance

- Ingredients

- Active ingredients

Acriflavine

- Inactive ingredients

Deionised Water, Sodium Benzoate

- MAL number

MAL19910649XZ

Manufacturer & Product registration holder

Dynapharm (M) Sdn. Bhd.

2497, Mk 1, Lorong Perusahaan Baru 5,

Kawasan Perusahaan Perai 3, 13600

Perai, Pulau Pinang, Malaysia.

Date of revision

06/11/2024

Serial Number:

NPRA(R4/1)251018/00302