

Outer Label For Plain Box of Blister Pack of 100x10's:This Design Also Applicable For Blister Pack of 1x10's and 10x10's

**CONTROLLED
MEDICINE**

LB1PT D035-05-H26

After assessing the risk/benefit ratio in each individual patient, the lowest effective dose for the shortest possible duration should be used.

DOSAGE :

Adult : Pain: 500 mg initially, followed by 250 mg every 6 hours as necessary. Treatment should not be continued for longer than 7 days. Children with Still's disease : 25mg/kg bodyweight daily in divided doses. Dysmenorrhoea : medication should be initiated with the onset of bleeding and associated symptoms and should not be necessary for more than 2-3 days. All to be taken with some food or after meal.

INDICATIONS :

Mild to moderate pain such as muscular aches, backaches, dental pain, bursitis, dysmenorrhoea, rheumatic pain, traumatic pain, headaches of most aetiology, postoperative and postpartum pain, rheumatoid arthritis (including Still's disease), osteoarthritis and menorrhagia due to dysfunctional causes.

Dyfenamic Tablet **500 mg** (film-coated)


Each tablet contains :
Mefenamic Acid 500 mg

Batch Size :
Mfg Date :
Exp Date :

Blister pack of 100x 10's MAL19910166AZ



Scan QR for Package Insert




9 557490 804355

JAUHI UBAT DARIPADA KANAK-KANAK.
KEEP OUT OF REACH OF CHILDREN.
PROTECT FROM LIGHT. DO NOT STORE ABOVE 30°C.
FOR ORAL USE ONLY

Manufacturer / Product Registration Holder:
DYNAPHARM (M) SDN. BHD. 198001011897 (65683-V)
2497, Mk. 1, Lorong Perusahaan Baru 5,
Kawasan Perusahaan Perai 3,
13600 Perai, Pulau Pinang, MALAYSIA.



Plain Box For Blister Pack of 100x10's:



Note:

The Outer Label For
Plain Box of Blister Pack of 100 x 10's
(As At Pg. 1 of 2)
Shall Be Pasted Herewith.