

DYNA AMOXYCILLIN CAPSULE

Amoxicillin Trihydrate (250 mg, 500 mg)

What is in this leaflet

1. What DYNA AMOXYCILLIN CAPSULE is used for
2. How DYNA AMOXYCILLIN CAPSULE works
3. Before you use DYNA AMOXYCILLIN CAPSULE
4. How to use DYNA AMOXYCILLIN CAPSULE
5. While you are using it
6. Side effects
7. Storage and Disposal of DYNA AMOXYCILLIN CAPSULE
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What DYNA AMOXYCILLIN CAPSULE is used for

DYNA AMOXYCILLIN CAPSULE is used for

DYNA AMOXYCILLIN CAPSULE is used for

- Infections caused by bacteria
- Ear nose and throat infections
- Chest infections
- Urinary tract infections
- Skin and soft tissue infections

How DYNA AMOXYCILLIN CAPSULE works

AMOXYCILLIN CAPSULE works

Amoxicillin work by killing the bacteria that cause infections.

Before you use DYNA AMOXYCILLIN CAPSULE

- When you must not use it

Not to be use in individuals with known hypersensitivity to penicillin.

Pregnancy and lactation

Do not take DYNA AMOXYCILLIN CAPSULE if you are pregnant, trying to get pregnant or think you may be pregnant.

Do not take DYNA AMOXYCILLIN CAPSULE if you are breast-feeding. Ask your doctor or pharmacist for advice before taking any medicine.

- Before you start use it

Tell your doctor if:

- You are hypersensitive to other penicillins or cephalosporins or penicillamine.
- Impaired kidney function

Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

Probenecid prolongs blood levels of amoxicillin. Antacids and other alkalisng agents can impair absorption of amoxicillin. May reduce effectiveness of oral contraceptives. Concurrent administration of amoxicillin and allopurinol increases incidence of rashes.

How to use DYNA AMOXYCILLIN CAPSULE

- How much to use

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

- When to use it

Use as directed by your doctor or pharmacist.

- How long to use it

Continue taking DYNA AMOXYCILLIN CAPSULE for as long as your doctor recommends.

- If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too many tablets may cause gastrointestinal system disturbances, reversible impairment of liver function and diarrhoea.

While you are using it

- Things you must do

Take your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are taking DYNA AMOXYCILLIN CAPSULE.

Tell your doctor immediately if you become pregnant while taking this medication.

- Things you must not do

Do not stop taking the medicine unless advised by your doctor.

Do not take any new medicines without consulting your doctor.

Do not give DYNA AMOXYCILLIN CAPSULE to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

- You should continue medicine for full course of treatment.
- Take medicine on empty stomach.
- Reduced dosage is required in patients with impaired kidney function.

Side effects

Like all medicines, DYNA

AMOXYCILLIN CAPSULE can cause side effects, although not everybody gets them.

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine.

Side effects are usually mild and transitory e.g. nausea, vomiting, diarrhoea, indigestion. Amoxicillin may give rise to a rash which may be of toxic origin. Rashes almost invariably result when Amoxicillin is given to individuals with glandular fever (viral infection). Small amount of Amoxicillin excreted in the milk may provoke allergic reactions in breast-fed infants.

Stop taking DYNA AMOXYCILLIN CAPSULE and contact your doctor immediately if you experience any of the following:

- Serious allergic reactions such as Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS). DRESS

DYNA AMOXYCILLIN CAPSULE

Amoxicillin Trihydrate (250 mg, 500 mg)

appears initially as flu-like symptoms with a rash on the face and then with an extended rash, high temperature and enlarged lymph nodes.

DYNA AMOXYCILLIN CAPSULE may cause severe allergy and serious skin reactions.

Stop using DYNA AMOXYCILLIN CAPSULE and seek medical assistance immediately if you experience any of the following symptoms:

- Skin reddening, blisters, rash, fever, sore throat or eye irritations.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my (Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)).

Storage and Disposal of DYNA AMOXYCILLIN CAPSULE

- Storage

KEEP OUT OF REACH OF CHILDREN

Do not store above 30°C

Protect from light

Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

DYNA AMOXYCILLIN CAPSULE
250MG

Orange body/red cap hard gelatin capsule of size “2” with white to off-white granules.

DYNA AMOXYCILLIN CAPSULE
500MG

Orange body/red cap hard gelatin capsule of size “0” with white to off-white granules.

Ingredients

- Active ingredient(s)

Amoxicillin Trihydrate

- Inactive ingredients

DYNA AMOXYCILLIN CAPSULE

250MG

Lactose powder 200 mesh, Magnesium Stearate.

DYNA AMOXYCILLIN CAPSULE

500MG

Magnesium Stearate

MAL number(s):

DYNA AMOXYCILLIN CAPSULE

250MG

MAL19930222AZ

DYNA AMOXYCILLIN CAPSULE

500MG

MAL19930221AZ

Manufacturer and Product

Registration Holder

Dynapharm (M) Sdn. Bhd.

2497, Mk 1, Lorong Perusahaan Baru 5,

Kawasan Perusahaan Perai 3, 13600

Perai, Pulau Pinang, Malaysia.

Date of revision

08/04/2022

Serial Number

NPRA(R2/1)201223/587