

The Outer Box For **Aluminium Tube of 10 g:**

		 JAUHI UBAT DARIPADA KANAK-KANAK KEEP OUT OF REACH OF CHILDREN Protect from light Store in dry place below 30°C Keep container tightly closed Please refer to package insert for further information.	
		 Manufacturer/Product Registration Holder: DYNAPHARM (M) SDN. BHD. 198001011897 (65683-V) 2497, MK1, Lorong Perusahaan Baru 5, Kawasan Perusahaan Perai 3, 13600 Perai, Pulau Pinang, Malaysia.  9 557490 604054	
	 DYNA ORAPASTE Treatment and relief of oral inflammatory lesions MAL19984055AZ GEL FOR ORAL APPLICATION ONLY CONTROLLED MEDICINE 10 grams Triamcinolone 0.1% w/w (as Acetonide) Preservatives: Methyl Paraben 0.2% w/w & Propyl Paraben 0.02% w/w	Batch No. : Mfg. Date : Exp. Date : BX1PE D005-07-I24	
	INDICATION: Indicated for adjunctive treatment and for temporary relief of symptoms associated with oral inflammatory lesions and ulcerative lesions resulting from trauma. DIRECTION FOR USE: Apply a small dab of DYNA ORAPASTE to the lesion. A large quantity may be required for coverage of some lesions. For optimal results use only enough to coat the lesion with a thin film. The medication should be applied at bedtime to permit steroid contact with the lesion throughout the night. Depending on the severity of symptoms, it may be necessary to apply the preparation 2 or 3 times a day, preferably after meals.		