

DYNA ORAPASTE

Triamcinolone Acetonide (0.1% w/w)

What is in this leaflet

1. What Dyna Orapaste is used for
2. How Dyna Orapaste works
3. Before you used Dyna Orapaste
4. How to use Dyna Orapaste
5. While you are using it
6. Side effects
7. Storage and disposal of Dyna Orapaste
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Dyna Orapaste is used for

Dyna Orapaste indicated for adjunctive treatment and for temporary relief of symptoms associated oral inflammatory lesion and ulcerative lesions resulting from trauma.

How Dyna Orapaste works

Triamcinolone provides a synthetic corticosteroid with mark anti-inflammatory action. It also provides a soothing and protective covering, which augments the effects of the steroid.

Before you use Dyna Orapaste

- When you must not use it

Do not use Dyna Orapaste if you have known hypersensitivity to Triamcinolone, corticosteroids, propylene glycol or other ingredients listed in this leaflet.

Pregnancy and lactation

Tell your doctor if you are pregnant, think you may be pregnant or planning to get pregnant or if you are breastfeeding. Caution should be exercised when used by nursing mothers.

Infants born of the mothers who have received substantial doses of corticosteroids during pregnancy should be carefully observed for signs of underactive adrenal glands such as hypoadrenalism (weight loss, dizziness, stomachache, patches of darker skin).

- Before you start to use it

Tell your doctor or pharmacist if you have any of the following conditions:

- Tuberculosis (bacterial infection that mainly affects the lungs)
- Stomach or intestinal ulcer
- Diabetes
- Viral infection

- Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescriptions from a pharmacy, supermarket or health food shop.

How to use Dyna Orapaste

- How much to use

Follow all directions given to you by your doctor and pharmacists carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

Apply a small dab of Triamcinolone-containing gel to the lesion. A larger quantity may be required for coverage of some lesions. For optimal results use only enough to coat the lesion with a thin film.

WARNING

Discontinue if local irritations develop.

- When to use it

Use as directed by your doctor or pharmacist.

The gel should be applied at bedtime to permit steroid contact with the lesion throughout the night. Depending on the severity of symptoms, it may be necessary to apply the preparation 2 or 3 times a day, preferably after meals.

- How long to use it

Continue using Dyna Orapaste for as long as your doctor recommends. Consult your doctor in 7 days if your condition did not improve.

- If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

- If you use too much (overdose)

This gel is for external use, there are no symptoms of an acute overdose known

While you are using it

- Things you must do

Use your medicine exactly as your doctor has told you.

Tell the doctor, dentist and pharmacist treating you that you are using Dyna Orapaste

- Things you must not do

Do not use any new medicines without consulting your doctor or pharmacist.

Do not give Dyna Orapaste to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

Immunisation procedures should not be undertaken in people who are on corticosteroids, especially on high dose, because of possible hazards of neurological complications and lack of antibody response.

Side effects

Like all medicines, Dyna Orapaste can cause side effects, although not everybody gets them.

Intolerance to Triamcinolone – containing gel is extremely rare. However, if any unusual symptoms, such as weakness or dizziness should occur, you should inform your doctor.

Visit your doctor or pharmacist immediately if you experience any side effects after using this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my (Consumers → Reporting → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)).

Storage and Disposal of Dyna Orapaste

- Storage

Keep out of reach of children
Store in dry place below 30°C
Keep the tube tightly closed

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

DYNA ORAPASTE

Triamcinolone Acetonide (0.1% w/w)

Product Description

- What it looks like

Natural (transparent) in colour gel
with apricot flavour

- Ingredients

- Active ingredients

Triamcinolone Acetonide

- Inactive ingredients

Purified Water, Methyl Paraben,
Propyl Paraben, Citric Acid
Anhydrous, Methocel E15LV (Cotacel
C15/HPMC), Carboxymethylcellulose
Sodium, Glycerin & Apricot Flavour.

- MAL number

MAL19984055AZ

Manufacturer & Product

Registration Holder

Dynapharm (M) Sdn. Bhd.
2497, Mk 1, Lorong Perusahaan Baru
5, Kawasan Perusahaan Perai 3,
13600 Perai, Pulau Pinang, Malaysia.

Date of revision

26/09/2024

Serial Number:

NPRA(R4/1)141217/00360