

DYNA AMOXYCILLIN SUSPENSION

Amoxicillin Trihydrate (125 mg/5 ml, 250 mg/5 ml)

What is in this leaflet

1. What DYNA AMOXYCILLIN is used for
2. How DYNA AMOXYCILLIN works
3. Before you use DYNA AMOXYCILLIN
4. How to use DYNA AMOXYCILLIN
5. While you are using it
6. Side effects
7. Storage and Disposal of DYNA AMOXYCILLIN
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What DYNA AMOXYCILLIN is used for

DYNA AMOXYCILLIN is used for

- Infections caused by bacteria.
- Ear nose and throat infections
- Chest infections.
- Urinary tract infections
- Skin and soft tissue infections.

How DYNA AMOXYCILLIN works

Amoxicillin work by killing the bacteria that cause infections.

Before you use DYNA AMOXYCILLIN

- When you must not use it

Not to be used in patients with known hypersensitivity to penicillin.

Pregnancy and lactation

Do not take DYNA AMOXYCILLIN if you are pregnant, trying to get pregnant or think you may be pregnant.

Do not take DYNA AMOXYCILLIN if you are breast-feeding. Ask your doctor or pharmacist for advice before taking any medicine.

- Before you start use it

Tell your doctor if:

- You are hypersensitive to other penicillins or cephalosporins or penicillamine.
- Impaired kidney function

- Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

- Probenecid prolongs blood levels of amoxicillin.

- Antacids and other alkalisating agents can impair absorption of amoxicillin.
- May reduce effectiveness of oral contraceptives.
- Concurrent administration of amoxicillin and allopurinol increases incidence of rashes.

How to use DYNA AMOXYCILLIN

- How much to use

Shake to loosen powder. Add approximately half the final volume (30 ml or 50 ml) of freshly boiled and cooled water and shake gently.

Add more water until the final volume (60 ml or 100 ml) mark is reached and shake well.

Keep refrigerated and to be used within 7 days of mixing.

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

- When to use it

Use as directed by your doctor or pharmacist.

- How long to use it

Continue taking DYNA AMOXYCILLIN for as long as your doctor recommends.

- If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Problem of overdosage are unlikely to be encountered. Gross overdosage will produce very high urinary concentration.

While you are using it

- Things you must do

Take your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are taking DYNA AMOXYCILLIN.

Tell your doctor immediately if you become pregnant while taking this medication.

- Things you must not do

Do not stop taking the medicine unless advised by your doctor.

Do not take any new medicines without consulting your doctor.

Do not give DYNA AMOXYCILLIN to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

- You should continue medicine for full course of treatment.
- Take medicine on empty stomach.
- Reduced dosage is required in patients with impaired kidney function.

Side effects

Like all medicines, DYNA AMOXYCILLIN can cause side effects, although not everybody gets them.

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine.

Side effects are usually mild and transitory e.g. nausea, vomiting, diarrhoea, indigestion. Amoxicillin may give rise to a rash which may be of toxic origin. Rashes almost invariably result when Amoxicillin is given to patients with glandular (viral infection) fever. Small amount of Amoxicillin excreted in the milk may provoke allergic reactions in breast-fed infants. Inflammation of the colon.

Stop taking DYNA AMOXYCILLIN and contact your doctor immediately if you experience any of the following:

- Serious allergic reactions such as Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS). DRESS appears initially as flu like symptoms with a rash on the face and then with an

DYNA AMOXYCILLIN SUSPENSION

Amoxicillin Trihydrate (125 mg/5 ml, 250 mg/5 ml)

extended rash, high temperature and enlarged lymph nodes.

DYNA AMOXYCILLIN may cause severe allergy and serious skin reactions.

Stop using DYNA AMOXYCILLIN and seek medical assistance immediately if you experience any of the following symptoms:

- Skin reddening, blisters, rash, fever, sore throat or eye irritations.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my (Consumers → Reporting → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)).

Storage and Disposal

- Storage

DYNA AMOXYCILLIN 125MG/5ML

Keep out of reach of children

Keep container tightly closed

Do not store above 30°C

Protect from light.

DYNA AMOXYCILLIN 250MG/5ML

Keep out of reach of children

Keep container tightly closed

Do not store above 30°C

Protect from light.

- Storage (after reconstitution)

Keep refrigerated (2-8°C) after reconstitution

To be used within 7 days of mixing

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

DYNA AMOXYCILLIN 125MG/5ML

White coloured powder. Pink in colour suspension, opaque and smooth with strawberry flavour on adding water.

DYNA AMOXYCILLIN 250MG/5ML

White coloured powder. Pink in colour suspension, opaque and smooth with strawberry flavour on adding water.

- Ingredients

- Active ingredient(s)

Amoxicillin Trihydrate

- Inactive ingredients

Aerosil 200, Strawberry Flavour Powder, Red Carmoisine, Rhodigel (Xanthan Gum), Sugar (Crushed Fine), Sodium Lauryl Sulphate, Sodium Benzoate, Sodium Saccharin.

- MAL number(s):

DYNA AMOXYCILLIN 125MG/5ML

MAL19985981AZ

DYNA AMOXYCILLIN 250MG/5ML

MAL19985982AZ

Manufacturer and Product

Registration Holder

Dynapharm (M) Sdn. Bhd.

2497, Mk 1, Lorong Perusahaan Baru 5,

Kawasan Perusahaan Perai 3,

13600 Perai, Pulau Pinang, Malaysia.

Date of revision

02/11/2023

Serial Number

NPRA(R4/1)210917/00254