

Immediate Label For Plastic Bottle of 60 ml:This Design Also Applicable For Plastic Bottles of 100 ml**Direction for Mixing :**

Shake to loosen powder.
Add approximately 30ml
freshly boiled and cooled
water and shake well.
Add more water until 60ml
mark is reached and shake.

Keep refrigerated (2-8°C)
after reconstitution. To be
used within 7 days of mixing.

Complete the prescribed
course.

**NOT TO BE USED IN
PATIENTS WITH KNOWN
HYPERSENSITIVITY TO
PENICILLIN**



Manufacturer / Product Registration
Holder:

DYNAPHARM (M) SDN. BHD.

198001011897 (65683-V)

2497, Mk 1, Lorong Perusahaan Baru 5,
Kawasan Perusahaan Perai 3,
13600 Perai, Pulau Pinang, MALAYSIA.

Dyna Ampicillin

125mg / 5ml

**Each 5 ml of the reconstituted
suspension contains :-
Ampicillin Trihydrate
equivalent to Ampicillin 125 mg
Preservative :**

Sodium Benzoate.....0.1% w/v

FOR ORAL USE ONLY

60 ML MAL19985980AZ

Indications and Dosage :

Refer To Package Insert.

SHAKE WELL BEFORE USE

JAUHI UBAT DARIPADA KANAK-KANAK.

KEEP OUT OF REACH OF CHILDREN.

DO NOT STORE ABOVE 30°C

PROTECT FROM LIGHT.

KEEP CONTAINER TIGHTLY CLOSED.

CONTROLLED MEDICINE

Batch No.:

Mfg. Date:

Exp. Date:

LB1PL A007-10-J23

