

# DYNA CLOXACILLIN CAPSULE

Cloxacillin Sodium (250mg, 500mg)

## What is in this leaflet

1. What Dyna Cloxacillin Capsule is used for
2. How Dyna Cloxacillin Capsule works
3. Before you use Dyna Cloxacillin Capsule
4. How to use Dyna Cloxacillin Capsule
5. While you are using it
6. Side effects
7. Storage and Disposal of Dyna Cloxacillin Capsule
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

## What Dyna Cloxacillin Capsule is used for

Dyna Cloxacillin Capsule is used to treat infections caused by resistant bacteria, including

- Infections of skin and soft tissues
- Bones and joints
- Respiratory tract and urinary tract
- Otitis media
- Inflammation on the heart's inner lining
- Blood poisoning
- Inflammation of the membranes surrounding brain and spinal cord.

## How Dyna Cloxacillin Capsule works

Cloxacillin is a bactericidal antibiotic which inhibits bacterial cell wall synthesis.

## Before you use Dyna Cloxacillin Capsule

### - When you must not use it

Do not take Dyna Cloxacillin Capsule if you have hypersensitive to Cloxacillin or a history of allergic to Penicillin group of drugs.

If a skin rash occurs, treatment should be discontinued.

### *Pregnancy and lactation*

Do not take Dyna Cloxacillin Capsule if you are pregnant, trying to get pregnant or think you may be pregnant.

Do not take Dyna Cloxacillin Capsule if you are breast-feeding. Ask your doctor or pharmacist for advice before taking any medicine.

### - Before you start use it

Tell your doctor if you have impaired kidney function, hypersensitive to any

other Penicillin or Cephalosporin or Penicillamine.

### - Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

Probenecid impairs kidney excretion and reduces anticoagulant effect.

## How to use Dyna Cloxacillin Capsule

### - How much to use

Adult: 250 – 500 mg every 6 hours daily.

Children: 50-100 mg per kg body-weight daily in divided doses

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

### - When to use it

Use as directed by your doctor or pharmacist.

Take medicine on an empty stomach.

### - How long to use it

Continue taking Dyna Cloxacillin Capsule for as long as your doctor recommends.

You or your child should continue medicine for full course of treatment.

### - If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

### - If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too many tablets may cause gastrointestinal disturbance, reversible impairment of liver function and diarrhea.

## While you are using it

### - Things you must do

Take your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are taking Dyna Cloxacillin Capsule.

Tell your doctor immediately if you are pregnant while taking this medication.

### - Things you must not do

Do not stop taking the medicine unless advised by your doctor.

Do not take any new medicines without consulting your doctor.

Do not give Dyna Cloxacillin Capsule to anyone else, even if they have the same symptoms or condition as you.

### - Things to be careful of

Tell your doctor / pharmacist if skin rash occur. Reduced dosage is required in individuals with impaired renal function.

## Side effects

Like all medicines, Dyna Cloxacillin Capsule can cause side effects, although not everybody gets them.

Side effects such as diarrhoea, nausea and heartburn may occur following administration by mouth. Sensitivity reactions may occur to individuals hypersensitive to the drug. The most common side effects are allergic reaction and of these, skin rashes occur most frequently.

Dyna Cloxacillin Capsule may cause severe allergy and serious skin reactions.

Stop using Dyna Cloxacillin Capsule and seek medical assistance immediately if you experience any of the following symptoms: skin reddening, blister, rash, fever, sore throat or eye irritation.

# DYNA CLOXACILLIN CAPSULE

Cloxacillin Sodium (250mg, 500mg)

---

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website [npra.gov.my](http://npra.gov.my) (Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)).

## Manufacturer and Product

### Registration Holder

Dynapharm (M) Sdn. Bhd.  
2497, Mk. 1, Lorong Perusahaan Baru  
5, Kawasan Perusahaan Perai 3,  
13600 Perai, Pulau Pinang, Malaysia.

### Date of revision

04/03/2025

### Serial Number:

NPRA(R4/1)240418/00120

## Storage and Disposal of Dyna Cloxacillin Capsule

### - Storage

Store in a dry place below 30°C  
Protect from light  
Keep out of reach of children

### - Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

## Product Description

### - What it looks like

Dyna Cloxacillin Capsule 250mg  
Orange/black hard gelatin capsule of size "2" with white granules.

Dyna Cloxacillin Capsule 500mg  
Amethyst/black hard gelatin capsule of size "0" with white granules and marking 'DYNA' and company logo.

### - Ingredients

- Active ingredient  
Cloxacillin Sodium

- Inactive ingredients  
Dyna Cloxacillin Capsule 250mg  
Lactose Powder 200 Mesh, Magnesium Stearate, Corn Starch

Dyna Cloxacillin Capsule 500mg  
Lactose Powder 200 Mesh, Magnesium Stearate

### - MAL number(s):

Dyna Cloxacillin Capsule 250mg  
MAL19930225AZ

Dyna Cloxacillin Capsule 500mg  
MAL07090901AZ