

VITBION FORTE TABLET

Thiamine HCl / Pyridoxine HCl / Cyanocobalamin (100 mg / 200 mg / 200 mcg)

What is in this leaflet

1. What VITBION FORTE TABLET is used for
2. How VITBION FORTE TABLET works
3. Before you used VITBION FORTE TABLET
4. How to use VITBION FORTE TABLET
5. While you are using it
6. Side effects
7. Storage and disposal of VITBION FORTE TABLET
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What VITBION FORTE TABLET is used for

It is used as a supplement in the treatment of inflammation involving nerves, irritated or damage nerves, diabetes and drugs induced damage nerves.

How VITBION FORTE TABLET works

- Thiamine is fundamentally associated with carbohydrate metabolism.
- Pyridoxine is the coenzyme for a variety of metabolic functions affecting protein, carbohydrate and lipid utilization.
- Cyanocobalamin is used in the treatment of anemia and nerves complication and also associated with nutritional deficiencies.

Before you use VITBION FORTE TABLET

- When you must not use it

Ask your doctor or pharmacists for advice before taking VITBION FORTE TABLET.

- Before you start to use it

Tell your doctor if you have hypersensitivity to the ingredients.

- Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

How to use VITBION FORTE TABLET

- How much to use

For therapeutic use: 1 tablet 3 times daily.

The directions given may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

- When to use it

Use as directed by your doctor or pharmacist.

- How long to use it

Continue taking VITBION FORTE TABLET for as long as your doctor recommends.

- If you forget to use it

Take the missed dose as soon as you remember. If it almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

Overdosage rarely occur. Excess Vitamin B will be excreted in the urine.

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using it

- Things you must do

Take your medicine exactly as your doctor has told you.

Tell the doctor, dentist and pharmacist treating you that you are taking VITBION FORTE TABLET.

- Things you must not do

Do not take any new medicines without consulting your doctor or pharmacist.

Do not give VITBION FORTE TABLET to anyone else, even if they have the same symptoms or condition as you.

Side effects

Vitamin B groups are water soluble vitamins and they do not cause any side effects if taken accordingly to the recommended dosage.

Warning:

This product contains Vitamin B6.

Stop taking this product if you experience tingling, burning or numbness and see your doctor as soon as possible.

Hypersensitivity reactions to Vitamin B1, such as sweating, tachycardia (rapid heartbeat), and skin reactions with itching and urticaria are very rare.

Chromaturia ("reddish urine", appears during the first 8 hours after an administration and typically resolves within 48 hours) may occur.

Gastrointestinal complaints, such as nausea, vomiting, diarrhoea or abdominal pain may occur.

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my (Public → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)).

Storage and Disposal of VITBION FORTE TABLET.

- Storage

Keep out of reach of children
Protect from light
Do not store above 30°C

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

Orange colour tablet with round, biconvex and film coated tablet.

- Ingredients

- Active ingredients
Thiamine HCl, Pyridoxine HCl, Cyanocobalamin

- Inactive ingredients

Colorcon Orange, Magnesium Stearate, Corn Starch, Lactose Powder 200 Mesh, PVP K90 & Aerosil 200.

VITBION FORTE TABLET

Thiamine HCl / Pyridoxine HCl / Cyanocobalamin (100 mg / 200 mg / 200 mcg)

- MAL number

MAL19920011XZ

Manufacturer & Product Registration

Holder

Dynapharm (M) Sdn. Bhd.
2497, Mk. 1, Lorong Perusahaan Baru 5,
Kawasan Perusahaan Perai 3,
13600 Perai, Pulau Pinang, Malaysia.

Date of Revision

02/07/2026

Serial Number:

NPRA(R4/1)170518/00156