

DYNA LAXATIVE PILL

Bisacodyl (5 mg)

What is in this leaflet

1. What Dyna Laxative Pill is used for
2. How Dyna Laxative Pill works
3. Before you used Dyna Laxative Pill
4. How to use Dyna Laxative Pill
5. While you are using it
6. Side effects
7. Storage and disposal of Dyna Laxative Pill
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Dyna Laxative Pill is used for

Indicated for the treatment of constipation and bowel evacuation before investigation procedures or surgery.

How Dyna Laxative Pill works

Bisacodyl is known as a stimulant laxative. It works by increasing the movement of the intestines, helping the stool to come out.

Before you use Dyna Laxative Pill

- When you must not use it

Dyna Laxative Pill should not be used if you have intestinal obstruction, abdominal pain, nausea or vomiting.

Pregnancy and lactation

Do not take Dyna Laxative Pill if you are pregnant, trying to get pregnant or think you may be pregnant especially in the first trimester. Do not take Dyna Laxative Pill if you are breast-feeding.

Ask your doctor or pharmacists for advice before taking Dyna Laxative Pill.

- Before you start to use it

Tell your doctor if you are hypersensitive to Bisacodyl and other ingredients listed in the leaflet.

- Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

The concomitant use of antacids and milk products may result in indigestion and gastric irritation.

Used of diuretics or corticosteroids may increase risk of salt imbalance when taken excessively.

How to use Dyna Laxative Pill

- How much to use

Children 6 years and above: 1 tablet
Adults: 2 tablets

The directions given may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

- When to use it

Use as directed by your doctor or pharmacist.

To be taken before bedtime. Do not chew the tablet. Swallow whole with a glass of water. Do not exceed the stated dose unless advised by physician. Milk and antacid should not be taken within 1 hour.

- How long to use it

Continue taking Dyna Laxative Pill for as long as your doctor recommends. Avoid prolonged use.

- If you forget to use it

Take the missed dose as soon as you remember. If it almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

If high doses are taken, diarrhoea, abdominal cramps and loss of potassium and other salts can occur and also increase sensitivity to cardiac glycoside (an organic compound).

Long term overdose may cause secondary hyperaldosteronism leading to excess aldosterone hormone production) and kidney stone.

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using it

- Things you must do

Take your medicine exactly as your doctor has told you.

Tell the doctor, dentist and pharmacist treating you that you are taking Dyna Laxative Pill.

- Things you must not do

Do not take any new medicines without consulting your doctor or pharmacist.

Do not give Dyna Laxative Pill to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

Avoid prolonged use.

Side effects

Bisacodyl can cause abdominal discomfort. Prolonged used can cause diarrhoea, muscular weakness and weight loss.

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npa.gov.my (Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)).

Storage and disposal of Dyna Laxative Pill

- Storage

KEEP OUT OF REACH OF CHILDREN

Store in dry place below 30°C
Protect from light

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

DYNA LAXATIVE PILL

Bisacodyl (5 mg)

Product Description

- What it looks like

Brown colour with round and biconvex shape, enteric coated tablet.

- Ingredients

- Active ingredients

Bisacodyl

- Inactive ingredients

Corn Starch

Lactose Powder 200 Mesh

Sodium Starch Glycolate

PVP K90

Magnesium Stearate

Aerosil 200

Croscarmellose Sodium

PEG 4000

Methocel E15LV

Acryl-Eze Orange

- MAL number

MAL20014531XZ

Manufacturer & Product

Registration Holder

Dynapharm (M) Sdn. Bhd.

2497, Mk 1,

Lorong Perusahaan Baru 5,

Kawasan Perusahaan Perai 3,

13600 Perai, Pulau Pinang, Malaysia.

Date of Revision

01/03/2022

Serial Number:

NPRA(R2/1)220301/842