

DYNA PARACETAMOL SUSPENSION

Paracetamol (250 mg/ 5ml)

What is in this leaflet

1. What Dyna Paracetamol Suspension is used for
2. How Dyna Paracetamol Suspension works
3. Before you use Dyna Paracetamol Suspension
4. How to use Dyna Paracetamol Suspension
5. While you are using it
6. Side effects
7. Storage and disposal of Dyna Paracetamol Suspension
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Dyna Paracetamol Suspension is used for

Indicated for relief of fever and pain (such as headache and toothache) in children.

How Dyna Paracetamol Suspension works

Paracetamol acts by inhibiting the production of hormone-like substances that participate in a wide range of body functions such as the contraction and relaxation of muscle, the dilation and constriction of blood vessels, control of blood pressure and modulation of inflammation. It is considered to affect the brain that controls body temperature.

Before you use Dyna Paracetamol Suspension

- When you must not use it

Tell your doctor if you are hypersensitive to Paracetamol.

- Before you start to use it

Pregnancy and lactation

Pregnancy: Not known to be harmful

Lactation: Amount too small to be harmful.

Ask your doctor or pharmacists for advice before taking Dyna Paracetamol Suspension.

- Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

Paracetamol should not be taken concomitantly with

- Alcohol
- Cholestyramine (reduce cholesterol)
- Warfarin (prevent blood clot)
- Metoclopramide or Domperidone (treat nausea and vomiting)

Paracetamol should not be taken concomitantly with a medicine that acts on the liver.

How to use Dyna Paracetamol Suspension

- How much to use

Dyna Paracetamol Suspension (Pink)

Children aged 1 - 5 years: ½ to 1 teaspoonful (2.5 ml to 5 ml).

6 - 12 years: 1 to 2 teaspoonfuls (5 ml to 10 ml).

Doses may be repeated every 4 – 6 hours when necessary (maximum of 4 doses in 24 hours).

Dyna Paracetamol Suspension

Children 1 to 6 years: ½ to 1 teaspoonful

Children 7 to 12 years: 1 to 2 teaspoonfuls

To be given 3 to 4 times daily (1 teaspoonful = 5 ml)

The directions given may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

- When to use it

Use as directed by your doctor or pharmacist.

- How long to use it

Continue taking Dyna Paracetamol Suspension for as long as your doctor recommends.

Do not exceed the recommended dosage or use for more than 10 days unless prescribed.

If you forget to use it

Take the missed dose as soon as you remember. If it almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

Features of Paracetamol poisoning are nausea and vomiting, followed by pain and tenderness in the right side of lower ribs area of the abdomen. Overdose can cause liver damage that is signified by yellowing of the skin and eyes, gastrointestinal bleeding and low blood glucose level.

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using it

- Things you must do

Take your medicine exactly as your doctor has told you.

Tell the doctor, dentist and pharmacist treating you that you are taking Dyna Paracetamol Suspension.

- Things you must not do

Do not take any new medicines without consulting your doctor or pharmacist.

Do not give Dyna Paracetamol Suspension to anyone else, even if they have the same symptoms or condition as you.

This preparation contains PARACETAMOL. Do not take any other PARACETAMOL-containing medicines at the same time.

- Things to be careful of

This medicine should be used with care in liver and kidney impairment.

Side effects

Side effects of the Dyna Paracetamol Suspension usually mild. Nausea, gastro-intestinal upsets, rashes and allergic conditions have occurred.

DYNA PARACETAMOL SUSPENSION

Paracetamol (250 mg/ 5ml)

Allergy alert: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine. You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my (Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)).

Storage and Disposal of Dyna Paracetamol Suspension

Keep out of reach of children
Protect from light
Keep the container tightly closed
Shake well before use

Dyna Paracetamol Suspension (Pink)
Do not store above 30°C

Dyna Paracetamol Suspension
Store in dry place below 30°C

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

Dyna Paracetamol Suspension (Pink)
Pink colour suspension with raspberry flavour and fragrance.

Dyna Paracetamol Suspension
Yellow colour suspension with blended milk and vanilla flavour.

- Ingredients

- Active ingredients
Paracetamol

- Inactive ingredients

Dyna Paracetamol Suspension (Pink)
Raspberry Flavour, Red (Carmoisine), Propyl Paraben, Sorbitol 70% - Non Crystal, Xanthan Gum, Gum Tragacanth, Methyl Paraben, Sodium Saccharin, Sucrose (Refined Sugar) & Purified Water.

Dyna Paracetamol Suspension
Milk Flavour, Methyl Paraben, Propyl Paraben, Gum Tragacanth, Sodium Saccharin, Xanthan Gum, Syrup, Quinoline Yellow, Deionised Water, Sorbitol Solution and Vanilla Flavour.

- MAL number

Dyna Paracetamol Suspension (Pink)
MAL06091363XZ

Dyna Paracetamol Suspension
MAL19940225XZ

Manufacturer & Product Registration Holder

Dynapharm (M) Sdn. Bhd.
2497, Mk 1, Lorong Perusahaan Baru 5,
Kawasan Perusahaan Perai 3, 13600
Perai, Pulau Pinang, Malaysia.

Date of revision

31/01/2024

Serial Number:

NPRA(R2/2) 240116/1195