

# DYSOLVON SYRUP (RASPBERRY)

Bromhexine Hydrochloride (4 mg/5 ml)

## What is in this leaflet

1. What Dysolvon Syrup (Raspberry) is used for
2. How Dysolvon Syrup (Raspberry) works
3. Before you used Dysolvon Syrup (Raspberry)
4. How to use Dysolvon Syrup (Raspberry)
5. While you are using it
6. Side effects
7. Storage and disposal of Dysolvon Syrup (Raspberry)
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

## What Dysolvon Syrup (Raspberry) is used for

Administered in inflammation of the airway and other respiratory conditions to aid removal of phlegm or mucus.

## How Dysolvon Syrup (Raspberry) works

Bromhexine Hydrochloride has been reported to change the structure of airway secretions, increase the volume and reduce viscosity of the phlegm or mucus.

## Before you use Dysolvon Syrup (Raspberry)

### - When you must not use it

You are hypersensitive to Bromhexine or other ingredients of the formulation.

### - Before you start to use it

#### Pregnancy and lactation

Caution regarding drug usage during pregnancy should be practiced. Bromhexine should be avoided during lactation.

Ask your doctor or pharmacists for advice before taking Dysolvon Syrup (Raspberry)

### - Taking other medicines

Tell your doctor if you are taking antibiotics (Amoxicillin, Cefuroxime, Erythromycin or Doxycycline) and any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

## How to use Dysolvon Syrup (Raspberry)

### - How much to use

Adults: 2 - 4 teaspoonfuls (10 ml - 20 ml), 3 times daily.

Children under 5 years: 1 teaspoonful (5 ml), twice daily.

Children 5 to 10 years: 1 teaspoonful (5 ml), four time daily.

The directions given may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

### - When to use it

Use as directed by your doctor or pharmacist.

### - How long to use it

Continue taking Dysolvon Syrup (Raspberry) for as long as your doctor recommends.

### - If you forget to use it

Take the missed dose as soon as you remember. If it almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

### - If you use too much (overdose)

No reports regarding overdose.

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

## While you are using it

### - Things you must do

Take your medicine exactly as your doctor has told you.

Tell the doctor, dentist and pharmacist treating you that you are taking Dysolvon Syrup (Raspberry)

### - Things you must not do

Do not take any new medicines without consulting your doctor or pharmacist.

Do not give Dysolvon Syrup (Raspberry) to anyone else, even if they have the same symptoms or condition as you.

### - Things to be careful of

Bromhexine should be given cautiously to people with gastric ulceration.

## Side effects

Gastrointestinal side effects may occur occasionally a transient rise in liver enzyme transaminase in the blood has been reported.

Dysolvon Syrup (Raspberry) may cause severe allergy and serious skin reactions. Stop using this medicine and seek medical assistance immediately if you experience any of the following symptoms:

- 1) Severe allergy: breathing difficulties, light headedness, skin swellings or rash.
- 2) Severe skin reaction: skin reddening, blisters, rash, fever, sore throat or eye irritation.

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website [npra.gov.my](http://npra.gov.my) (Consumers → Reporting → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)).

## Storage and Disposal of Dysolvon Syrup (Raspberry)

### - Storage

Keep out of reach of children  
Protect from light  
Do not store above 30°C  
Keep container tightly closed

### - Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

# DYSOLVON SYRUP (RASPBERRY)

Bromhexine Hydrochloride (4 mg/5 ml)

---

## Product Description

- *What it looks like*

Colourless, raspberry flavour syrup

- *Ingredients*

- Active ingredients

Bromhexine Hydrochloride

- Inactive ingredients

Propylene Glycol, Raspberry Flavour,  
Menthol Crystals, Purified Water,  
Sugar, Citric Acid Anhydrous,  
Rhodigel (Xanthan Gum) & Sodium  
Benzoate.

- *MAL number*

MAL07091035XZ

## Manufacturer & Product

### Registration Holder

Dynapharm (M) Sdn. Bhd.  
2497, Mk 1, Lorong Perusahaan Baru  
5, Kawasan Perusahaan Perai 3,  
13600 Perai, Pulau Pinang, Malaysia.

## Date of Revision

05/03/2025

## Serial Number:

NPRA(R4/1)251018/00307