

CHILDREN'S PROTEMP SUPPOSITORIES 120MG

Paracetamol (120mg)

What is in this leaflet

1. What Children's Protemp Suppositories 120mg is used for
2. How Children's Protemp Suppositories 120mg works
3. Before you used Children's Protemp Suppositories 120mg
4. How to use Children's Protemp Suppositories 120mg
5. While you are using it
6. Side effects
7. Storage and disposal of Children's Protemp Suppositories 120mg
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Children's Protemp Suppositories 120mg is used for

Indicated for the temporary relief of fever, minor aches, pains and headaches.

How Children's Protemp Suppositories 120mg works

Paracetamol may acts by inhibiting prostaglandin synthesis in the central nervous system (CNS) and through a peripheral action by blocking pain-impulse generation. It also produce antipyretic effect by acting centrally on the hypothalamic heat regulating center to produce peripheral vasodilation resulting in increased blood flow through the skin and sweating and heat loss.

Before you use Children's Protemp Suppositories 120mg

- When you must not use it

Do not use Children's Protemp Suppositories 120mg to people with known hypersensitivity to Paracetamol and severe hepatic disease.

Pregnancy and lactation

Paracetamol crosses the placenta and distributes into breast milk.

Ask your doctor or pharmacists for advice before taking Children's Protemp Suppositories 120mg.

- Before you start to use it

Tell your doctor if you are patient of renal or with hepatic disease.

- Taking other medicines

- Patient who has taken barbiturates, tricyclic antidepressant and alcohol may risk of hepatotoxicity with prolonged use of high dose of Paracetamol because of increased metabolism resulting from induction of hepatic microsomal enzyme activity.
- Alcohol and hepatotoxic medication can increase the hepatotoxicity of Paracetamol overdose.
- Concurrent use anticoagulant, coumarin or indandione-derivative with chronic, high dose of Paracetamol may increase the anticoagulant effect.
- Prolonged concurrent use of Paracetamol with NSAIDS and salicylate may increase the renal adverse effect.

How to use Children's Protemp Suppositories 120mg

- How much to use

Dosing Chart	
Age	Dose
under 3 years	Do not use unless directed by a doctor
3 to 6 years	Use 1 suppository every 4 to 6 hours (maximum of 5 doses in 24 hours)

1. Wash your hand and remove suppository from wrapper by firmly grasp each tab and slowly peel apart the suppository in a downward motion.
2. Lie down on your left side with the right knee bent (if left-handed, lie on the right side with the left knee bent). Push the suppository into the rectum with your finger. Suppositories should remain in the rectum for 15 to 30 minutes.
3. If the suppository is too soft, run cold water over it before removing the foil wrapper for about 1-2 minutes.

The directions given may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

- When to use it

Use as directed by your doctor or pharmacist.

- How long to use it

Continue taking Children's Protemp Suppositories 120mg for as long as your doctor recommends.

- If you forgot to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

Overdose symptoms may include hepatic necrosis, transient azotemia, renal tubular necrosis with acute toxicity, anemia and GI disturbances with chronic toxicity. Treatment: Acetylcysteine 140 mg/kg orally (loading) followed by 70 mg/kg every 4 hours for 17 doses or by IV N-acetyl cysteine 150 mg/kg infused over 60 minutes followed by an IV maintenance dose of 50 mg/kg infused over 4 hours, and then 100 mg/kg infused over 16 hours. Supportive care includes maintaining fluid and electrolyte balance.

While you are using it

- Things you must do

Take your medicine exactly as your doctor has told you.

Tell the doctor, dentist and pharmacist treating you that you are taking Children's Protemp Suppositories 120mg.

- Things you must not do

Do not take any new medicines without consulting your doctor or pharmacist.

Do not give Children's Protemp Suppositories 120mg to anyone else, even if they have the same symptoms or condition as you.

CHILDREN'S PROTEMP SUPPOSITORIES 120MG

Paracetamol (120mg)

- Things to be careful of

If use more than the usually recommended dosage, on the label or the leaflet, hepatotoxic may occur. Patients with renal or hepatic disease should consult a physician or a pharmacist before use this drug.

This preparation contains PARACETAMOL.

Do not take any other paracetamol containing medicines at the same time.

Allergy alert: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

Side effects

Side effects are significantly mild, though hematological reactions have been reports.

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine. You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835490, or visiting the website npra.moh.gov.my (Public → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)).

Storage and Disposal of Children's Protemp Suppositories 120mg

- Storage

- Keep medicines out of reach of children.
- Store below 30°C
- For rectal use only

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

White, bullet shape suppositories.

- Ingredients

- Active ingredients
Paracetamol
- Inactive ingredients
Hard Fat

- MAL number

MAL*****

Manufacturer:

Lerd Singh Pharmaceutical Fact., Ltd.,
Part.
922 Soi Sukhumvit 50, Sukhumvit
Road, Phrakhanong, Khlongtoei,
Bangkok 10260, Thailand

Distributor / Product Registration

Holder:

Dynapharm (M) Sdn. Bhd.
2497, Mk 1, Lorong Perusahaan Baru 5,
Kawasan Perusahaan Perai 3, 13600
Perai, Pulau Pinang, Malaysia.

Date of revision

20/05/2019

Serial Number:

NPRA (R2) 19/92