

POTRELEASE TR TABLET

Potassium chloride (600mg)

What in this Leaflet

- 1.What Potrelease TR Tablet is used for
- 2.How Potrelease TR Tablet works
- 3.Before you use Potrelease TR Tablet
- 4.How to take Potrelease TR Tablet
- 5.While you are using it
- 6.Side Effects
- 7.Storage and Disposal of Potrelease TR Tablet
- 8.Products Descriptions
- 9.Manufacturer and Product Registration Holder
- 10.Date of revision

What Potrelease TR Tablet is used for

For treatment and prevention of low level of potassium in blood.

How Potrelease TR Tablet works

This medication is used to replenish potassium in blood.

Before you use Potrelease TR Tablet

-When you must not use it

Do not use Potrelease TR tablet if you are/have:

- Severe kidney impairment (you may produce little or no urine).
- Untreated (Addison's disease (inadequate production of hormones by adrenal glands)
- Dehydrated
- Heat cramps
- High level of potassium in blood
- Intestinal obstruction, abnormal movement or ulceration
- Too much acid in the body

-Before you start to use it

Please consult your doctor or pharmacist if you are pregnant, planning for pregnancy or breast-feeding before using any medicine

-Taking other medicine

Talk to your doctor/pharmacist if you are taking any medicine including those you buy without a prescription from a pharmacy, supermarket or health food shop. Medications that may interfere with Potrelease TR tablet are:

- Medication for high blood pressure: Captopril

- Slow heart rate and insecticide poisoning medication: Atropine
- Calcium salts
- Heart medication: Digitalis glycoside
- Hormone therapy: Adrenocorticoids, Glucocorticoids, Corticotrophin (ACTH)
- Medication for high blood potassium level: sodium polystyrene sulfonate
- Laxatives
- Low-salt milk
- Potassium- containing medications
- Water pills: Amiloride, Spironolactone, Triamterene
- Irregular heartbeat medication: Quinidine
- Vitamin B12

How to use Potrelease TR Tablet

-How much to use

- Adults: Oral, 1 to 2 tablets, three times a day.
- Children: Use is not recommended.
- Swallow tablets whole with a full glass of water.
- Do not chew or crush the tablet.

-When to use it

Take three times a day with or immediately after food or use as directed by your doctor or pharmacist.

-How long to use it

Use Potrelease TR Tablet for the duration as your doctor recommends.

-If you forget to use it

Take the missed dose as soon as possible as you remember it. However if it is almost for the next dose ,skip the missed dose and continue your regular doing schedule. Do not take a double dose to make up for the missed dose.

-If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have use too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using it

-Things you must do

Use your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are using Potrelease TR Tablet.

Tell your doctor immediately if you become pregnant while using this medication

-Things you must not do

Do not stop using the medicine unless advised by your doctor.

Do not use any new medicines without consulting your doctor or pharmacist.

Do not give Potrelease TR Tablet to anyone else, even if they have the same symptoms or condition as you.

-Things to be careful of

If you notice any side effects, stop using this medicine and tell your doctor or pharmacist.

Side Effects

Like all medicines, Potrelease TR Tablet can cause side effects, although not everybody gets them.

Side effects of Potrelease TR tablet are rare in people with normal kidney function. Consult your doctor immediately if you have: Confusion; Irregular heartbeat; Numbness or tingling in hands, feet, or lips; shortness of breath or difficulty in breathing; unexplained anxiety; unusual tiredness or weakness; weakness or heaviness of legs. Consult your doctor if these symptoms continue or are bothersome you: Diarrhoea, nausea, stomach pain and vomiting.

These side effects occur more frequently when the medication is not taken with food.

POTRELEASE TR TABLET

Potassium chloride (600mg)

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Manufacturer and Product Registration

Holder

SM PHARMACEUTICALS SDN.BHD.
Lot 88,Sungai Petani Industrial Estate
08000,Sungai Petani
Kedah Darul Aman,Malaysia

Date of Revision:22/02/2023

Serial Number:

NPRA(R4/1)050123/00002

Storage and Disposal of Potrelease TR Tablet

-Storage

Store below 30°C in a well-closed container.

Keep all medicines out of reach of children.

-Disposal

Medicines should not be disposed of via wastewater or household waste.

Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Products Description

- What it looks like

Yellow round shaped with deep concave surface film coated tablet

-Ingredients:

- Active ingredient
Potassium chloride.

- Inactive ingredients
Cetostearyl alcohol,ethyl
cellulose,gelatin,magnesium
stearate,isopropyl alcohol,purified
talcum,color quinoline yellow,tween
80,acetone,beeswax and purified water.

- MAL Number: MAL19950545XZ