

PREGAL 150 MG CAPSULE

Pregabalin 150 mg

What is in this leaflet

1. What Pregal 150mg capsule is used for

2. How Pregal 150mg capsule works

3. Before you take Pregal 150mg capsule

4. How to take Pregal 150mg capsule

5. While you are using Pregal 150 capsule

6. Side effects

7. Storage and Disposal of Pregal 150mg capsule

8. Product Description

9. Manufacturer and Product Registration Holder

10. Date of revision

What Pregal 150mg capsule is used for

- **Neuropathic pain**
Pregabalin is indicated for the treatment of peripheral and central neuropathic pain in adults.
- **Epilepsy**
Pregabalin is indicated as adjunctive therapy in adults with partial seizures with or without secondary generalisation.
- **Generalised Anxiety Disorder**
Pregabalin is indicated for the treatment of Generalised Anxiety Disorder (GAD) in adults.
- **Fibromyalgia**
Pregabalin is indicated for the management of fibromyalgia.

How Pregal 150mg capsule works

Pregabalin is in a class of medications called anticonvulsants. It works by decreasing the number of pain signals that are sent out by damaged nerves in the body.

Before you take Pregal 150mg capsule

- When you must not take it

If you are allergic to Pregabalin or any of the other ingredients of this medicine.

- Before you start to take it

Inform your doctor if you have

- Heart problem

- If you are pregnant or plan to get pregnant or breastfeeding.
- If you have diabetic

- Taking other medicines

Talk to your doctor / pharmacist if you are taking any medicine, including those you buy without prescription.

- Do not take Pregabalin if you are already taking other medicines containing Pregabalin.

- Do not take Pregal 150mg Capsule together with medicines containing

- **Other Antiepileptic** drugs (medications work in different ways to prevent seizures)
- **Oral Contraceptives** (are medications that prevent pregnancy.)
- **Lorazepam** (It is used to treat anxiety disorders)
- **Oxycodone** (is an opioid medication used for treatment of moderate to severe pain.)
- **Ethanol** (alcohols)
- **Diuretics** (are medications designed to increase the amount of water and salt expelled from the body as urine.)
- **Oral Hypoglycemics** (Drugs used in diabetes treat diabetes mellitus by lowering the glucose level in the blood)
- **Insulin** (is a hormone that lowers the level of glucose (a type of sugar) in the blood).

How to take Pregal 150mg Capsule

- How much to take

- The dose range is 150-600 mg/day given in either 2 or 3 divided doses.

- When to take it

Follow your doctor's advice

- How long to take it

Do not stop taking Pregal 150mg Capsule without your doctor's advice.

- If you forget to take it

It is important not to miss a dose of Pregal 150 capsule.

If you miss a dose, take it as soon as you remember it, and then take the next dose at its regular time.

If it is almost time for the next dose, skip the missed dose and continue your regular dosing schedule. Do not take a double dose to make up for a forgotten tablet.

- If you take too much (overdose)

If you have accidentally taken too many Celera capsule you may be at increased risk of experiencing possible side effects with this medicine.

Immediately contact your doctor or go to the nearest hospital for immediate medical attention.

While you are using Pregal 150 capsule

- Things you must do

Follow your doctor's advice.

- Things you must not do

Do not stop taking Pregal 150mg capsule without consulting your doctor first.

It is important to take precautions to avoid infecting other people.

Do not give your medicine to other people who experience the same symptoms as you.

- Things to be careful of

- Should be advised to seek medical advice should signs of suicidal ideation or behavior emerge.

PREGAL 150 MG CAPSULE

Pregabalin 150 mg

- Caution should be exercised in patients with a history of substance abuse and the patient should be monitored for symptoms of pregabalin misuse, abuse or dependence.

Ingredients

- Active ingredient used is Pregabalin
- Inactive ingredients: Pregelatinized starch
Purified Talc

Side effects

Like all medicines, this medicine can cause side effects although not everybody gets them. Possible serious side effects:

Common side effects include diarrhea, vomiting, nausea, dizziness.

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reactions Monitoring by visiting the website <http://nptra.moh.gov.my> Reporting Side Effects to Medicine (ConSERF) or Vaccines (Public(AEFI)).

MAL:

Manufacturer and Product Registration Holder

SM Pharmaceuticals Sdn. Bhd.
(218628-M)
Lot 88, Sungai Petani Industrial Estate
08000 Sungai Petani,
Kedah DarulAman, Malaysia

Date of revision: 30.08.2021
NPRA(R2)21/017

Storage and Disposal of Pregal 150mg Capsule

- Storage

Store below 30°C.

KEEP ALL MEDICINES OUT OF REACH OF CHILDREN.

- Disposal

Medicines should not be disposed of via waste water or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

Product Description

- What it looks like

Size '2' Capsules with white cap and white body imprinted with "SM-LU" on cap and "150" on body in black ink containing white to off white powder.