

Curekuf W Cough Syrup

Each 5ml contains Diphenhydramine Hydrochloride 14.0 mg and Ammonium Chloride 135.0 mg

What in this Leaflet

1. What is Curekuf W used for
2. How Curekuf W works
3. Before you use Curekuf W
4. How to take Curekuf W
5. While you are using it
6. Side Effects
7. Storage and Disposal of Curekuf W
8. Products Descriptions
9. Manufacturer and Product Registration Holder
10. Date of revision
11. Serial No.

What is Curekuf W used for

It is indicated as an expectorant for control of coughs due to colds or allergy.

How Curekuf W works

Diphenhydramine Hydrochloride is an antihistamine that works by blocking a certain natural substance (histamine) that your body makes during an allergic reaction.

Ammonium chloride acts by reflexing and irritating the gastric layer. These increases the volume of secretions in the respiratory tract; reduce thickness of phlegm which presumably is easier to cough up.

Before you use Curekuf W

-When you must not use it

Do not use Curekuf W if you

- are allergic to any ingredients in this medicine.
- not to be used in children less than 2 years of age.

-Before you start to use it

Tell your doctor or pharmacist before using Curekuf W if you:

- are pregnant or plan to get pregnant.
- are breast-feeding or plan to start breast-feeding.

-Taking other medicine

Tell your doctor or pharmacist if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop. Inform your doctor or pharmacist if you are taking:

- Alcohol
- Tranquilizers (a medicinal drug taken to reduce tension or anxiety).
- Sedative hypnotics (a class of drugs used to induce and/or maintain sleep).
- CNS depressant (medicines that include sedatives, tranquilizers, and hypnotics).
- MAO inhibitors (medication used to treat depression).

How to use Curekuf W

-How much to use

Adults and children over 12 years:

5 to 10 ml every 4 or 6 hours.

Do not to exceed 60 ml in 24 hours.

Children 2 to 5 years:

2.5 ml every 4 to 6 hours.

Do not to exceed 20 ml in 24 hours.

Children 6 to 12 years:

2.5 to 5 ml every 4 to 6 hours

Do not to exceed 30 ml in 24 hours.

-When to use it

Use as directed by your doctor or pharmacist.

-How long to use it

Continue using Curekuf W for the duration that the doctor recommends.

-If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

-If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have use too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using it

-Things you must do

Use your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are using Curekuf W.

Tell your doctor immediately if you become pregnant while using this medication. (if applicable)

Take Curekuf W only with a doctor's prescription. (For prescription only medicines)

-Things you must not do

Do not stop using the medicine unless advised by your doctor.

Do not use any new medicines without consulting your doctor or pharmacist.

Do not give Curekuf W to anyone else, even if they have the same symptoms or condition as you.

-Things to be careful of

Driving and using machines

This medicine may affect your ability to drive or use machines. If the syrup make you feel sick, dizzy or tired, or give you a headache, do not drive or use machines and contact your doctor immediately.

Curekuf W Cough Syrup

Each 5ml contains Diphenhydramine Hydrochloride 14.0 mg and Ammonium Chloride 135.0 mg

Side Effects

Like all medicines, Curekuf W can cause side effects, although not everybody gets them.

The most common side effects are sedation, sleepiness, dizziness, disturbed co-ordination, gastric irritation, and thickening of airway secretions. Less frequent side effects are hypersensitivity, skin rash, dry mouth, irritability, sleeping difficulties, euphoria (intense excitement and happiness), blurred vision, tinnitus (noise or ringing in the ears), spinning sensation, neuritis (inflammation of a peripheral nerve or nerves, usually causing pain and loss of function), and convulsion. Visit your doctor or pharmacist immediately if you experience any side effects after using this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring visiting the website npra.gov.my [Consumers→ Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of Curekuf W

-Storage

Shake well before use. Store in a dry place below 30°C. Protect from Light. Keep container tightly closed. Keep all medicines out of reach of children.

-Disposal

Medicines should not be disposed of via waste water or household waste. Ask your pharmacist how to dispose of the medicines no longer required. These measures will help you to protect the environment.

Products Description

-What it looks like

A brown colored, syrupy liquid with honey and pleasantly flavoured with peach and raspberry flavour.

-Ingredients:

-Active Ingredient used is Diphenhydramine hydrochloride and ammonium chloride

-Inactive ingredients used are Sodium citrate, sugar, menthol, maltodextrin, sodium benzoate, glycerin, sorbitol, propylene glycol, honey, liquid caramel, flavour raspberry, flavour peach and purified water

MAL No.:

MAL23026021AZ

-Packing

1x100ml per box

Manufacturer and Product

Registration Holder

SM PHARMACEUTICALS SDN.BHD
(218620-M)
Lot 88, Sungai Petani Industrial Estate
08000, Sungai Petani
Kedah Darul Aman, Malaysia

Date of Revision:

08/09/2023

Serial No.: NPRA (R2) 22/147