



STANDARDISED GINKGO BILOBA EXTRACT (EGb 761) – 40 mg

The medicinal product is a gastrosoluble coated tablet. It is round (diameter: approximately 9.1 mm), biconvex (radius of curvature: 12 mm), smooth and coloured a uniform red-brown. It is approximately 0.260 g in weight.

Composition (per tablet)

Dry extract from Ginkgo biloba (leaves/folium) (EGb 761) containing
22.0% - 27.0% of flavonoids expressed as flavone glycosides
5.4% - 6.6% of terpene lactones
2.8% - 3.4% of ginkgolides A, B, C
2.6% - 3.2% of bilobalide
Excipient q.s. one tablet

Therapeutic Indications

Traditionally used for improving blood circulation.

Adverse Effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Summary of the safety profile
The most common side effects are abdominal pain, diarrhea and dizziness.
If you develop a severe allergic reaction (angioedema) manifested by a swelling of your face, lips, tongue, or throat, stop taking this medicine immediately and seek medical attention as a matter of urgency.

- Other possible side effects:
Frequent (may affect up to 1 in 10 people)
- Allergic reaction
 - Difficulty in breathing
 - Headache
 - Syncope
 - Indigestion
 - Nausea
 - Eczema
 - Itching
- Uncommon (may affect up to 1 in 100 people)
- Hives
 - Rash

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

Dosage and Administration

Adult: The usual dosage is 1 tablet 3 times a day to be swallowed at mealtime.

Contra-indications/Drug-interactions

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

TANAKAN may alter the effect of other drugs: anticoagulants (phencoumarone, warfarin), anti-platelet drugs (clopidogrel, salicylic acid and other nonsteroidal anti-inflammatory drugs).

In case of concomitant use of this medicinal product with warfarin, monitoring is required, consult your doctor.

In case of concomitant use of this medicinal product with dabigatran etexilate, the effect of dabigatran may be potentiated, consult your doctor.

Concomitant use of this medication with efavirenz is not recommended due to the potential for decreased efavirenz effect.

Ginkgo can increase the concentration of nifedipine. In some individuals, this increase can be significant and cause dizziness and increased intensity of hot flashes.

		Customer	Mayoly Spindler Laboratories				Colours Used	Fonts Used	Notes
		SAP Code	7N00580ID				Keyline (Non-Printing)	Arial 6.5pt	
		SAP Description	NOT M TANAKAN MY				PANTONE 287 C	Helvetica Neue LT Std 7.5pt	
		Proof / Iteration	2				Text Free (Non-Printing)	Helvetica 7.1pt	
		Artwork Type	Leaflet				Cirrus_Info_Box		
Approval Signature		Profile Ref	N/A						
Date		Size	210 x 160mm						
Warning! We cannot accept responsibility for any errors in this proof after approval. Whilst we take extreme care at all times to ensure accuracy to our client's brief, the final responsibility must be taken by our client. IF YOU SIGN THIS PROOF YOU ARE SIGNIFYING FULL APPROVAL OF DESIGN AND TEXT.		Specification Ref	N/A						
		Barcode Type		Reel Direction	n 3				
		Pharmacode	67000580	Copy Position	N/A				
		Proof By	MDE	Date	19/08/2025				



This medicine is contraindicated during pregnancy.
There is insufficient reliable data in breastfeeding women.

Warning and Precaution

This is a traditional medicine.
Please consult your pharmacist/doctor before taking this product.
Keep out of reach of children. Jauhkan daripada capaian kanak-kanak.

The use of this drug is not recommended in patients with galactose intolerance, Lapp lactase deficiency or glucose or galactose malabsorption syndrome (rare hereditary diseases).

Concomitant use of this medicine with efavirenz is not recommended.

- Talk to your doctor before taking TANAKAN 40mg, coated tablet.
- If you have a predisposition to bleed (haemorrhagic predisposition) and if you have anticoagulant or anti-platelet therapy
 - If you suffer from epilepsy

Storage condition

Do not exceed the expiry date plainly indicated on the packaging.
Store below 30°C and protect from light and moisture.

Pack Size:

This medicine is in the form of coated tablets. Pack of 30 and 90 tablets

Product Registration Holder / Importer:
Zuellig Pharma Sdn Bhd
No. 15, Persiaran Pasak Bumi, Sek. U8,
Perindustrian Bukit Jelutong
40150, Shah Alam
Malaysia

Country of Origin: France
Date of revision: August 2025

Manufacturer: Mayoly Industrie – 20 rue Ethe Virton – 28100 Dreux – France

7N00580

		Customer	Mayoly Spindler Laboratories				Colours Used	Fonts Used	Notes
		SAP Code	7N00580ID				Keyline (Non-Printing)	Arial 6.5pt	
		SAP Description	NOT M TANAKAN MY				PANTONE 287 C	Helvetica Neue LT Std 7.5pt	
		Proof / Iteration	2				Text Free (Non-Printing)	Helvetica 7.1pt	
Approval Signature		Artwork Type	Leaflet				Cirrus_Info_Box		
		Profile Ref	N/A						
Date		Size	210 x 160mm						
		Specification Ref	N/A						
Barcode Type			Reel Direction	n 3					
Pharmacode		67000580	Copy Position	N/A					
Proof By		MDE	Date	19/08/2025					
Warning! We cannot accept responsibility for any errors in this proof after approval. Whilst we take extreme care at all times to ensure accuracy to our client's brief, the final responsibility must be taken by our client. IF YOU SIGN THIS PROOF YOU ARE SIGNIFYING FULL APPROVAL OF DESIGN AND TEXT.									