

BELLWYCK PACKAGING

FINAL PROOF

PLEASE CHECK CAREFULLY

ATTACHED IS A PROOF THAT NEEDS YOUR IMMEDIATE ATTENTION IN ORDER TO MEET YOUR REQUIRED DELIVERY SCHEDULE.

Customer: Bausch

Docket #: 138371

Proofer: Robert Davenport

Date: 2/2/2024

Proof # ☒ 1st ☐ 2nd ☐ 3rd ☐ ___ th

☐ Proof OK as is

☐ New Proof Required

ATTENTION - PLEASE READ THIS!

4 colour process (CMYK) areas are press-accurate. Pantone or custom spot colours are approximated and should NOT be used for colour evaluation. Changes made at this stage will be at your cost. Please proof carefully. What is shown in this proof represents what will be printed. We will not be responsible for errors if work is printed according to Customer's approval.

PLEASE FAX APPROVAL ASAP TO YOUR IN-HOUSE CSR AT (519) 376-8726.

APPROVAL DELAYS WILL RESULT IN A RE-SCHEDULING OF YOUR DELIVERY DATE

Customer Please Sign _____

Date _____

LZEBEL#



Proctosedyl® Ointment

Cinchocaine Hydrochloride, Hydrocortisone, Framycetin Sulphate, Aesculin

Rectal use

1 Tube (15g)

PHOENIX LABS

Manufacturer: Bausch Health Companies Inc., 2150 St. Elzear Blvd. West, Laval, Quebec, H7L 4A8, Canada.

Product Registration Holder & Marketed by: ZUELLIG PHARMA SDN BHD

No. 15, Persiaran Pasak Bumi, Sek. U8,
Perindustrian Bukit Jelutong,
40150 Shah Alam, Selangor Darul Ehsan,
Malaysia.

Proctosedyl® Ointment

Each gram contains:

Hydrocortisone 5mg

Cinchocaine hydrochloride 5mg

Framycetin sulphate 10mg

Aesculin 10mg

Other ingredients include woolfat (lanolin), liquid paraffin, white soft paraffin.

Controlled Medicine

Ubat Terkawal

MAL19871349ACZ

Keep out of reach of children. Jauhi daripada kanak-kanak.

Store below 30 °C (protect from excessive heat).

Discard contents 28 days after opening.

Indication: internal and external haemorrhoids; prophylaxis between attacks to prevent recurrences; haemorrhoidal complications - anal pruritis, perianal eczema, proctitis, fissure; pre-operative and postoperative treatment of haemorrhoidectomy patients; post-partum haemorrhoidal conditions.

Contraindications: Allergy to any of the ingredients, pregnancy or suspicion of pregnancy, breastfeeding or any existing infection.

Proctosedyl ointment should be applied around the back passage (or inside if needed) in the morning and at bedtime and, if possible, every time after bowels are emptied. Do not use for more than 7 days.

20005318A

LOT

MFG

EXP