

Calmoseptine® Ointment

Zinc Oxide(20.6%), Petrolatum(24%), Lanolin(15.7%), Menthol(0.44%)

What is in this leaflet

1. What Calmoseptine® Ointment is used for
2. How Calmoseptine® Ointment works
3. Before you use Calmoseptine® Ointment
4. How to use Calmoseptine® Ointment
5. While you are using it
6. Side effects
7. Storage and Disposal of Calmoseptine® Ointment
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of Revision.

What Calmoseptine® Ointment is used for

The ointment is used to temporarily protects irritated areas, temporary relief of local itching and discomfort in the perianal area, and for temporary relief of pain or burning or skin irritation.

How Calmoseptine® Ointment works

The ointment works as moisture barrier and astringent.

Before you use Calmoseptine® Ointment

-When you can't use the product

If you experience any allergy to the ingredients listed.

- Before you start to use it

Cleanse the affected area with mild soap and warm water and rinse thoroughly. Gently pat or air before application of this product.

- Taking other medicines

None known interaction with other medicines.

How to use Calmoseptine® Ointment

- How much to use

Adult: Apply externally to the affected area up to 6 times daily
Children under 12 years of age: Consult doctor before use.

- When to use it

Use up to 6 times or after each bowel movement.

- How long to use it

Apply until symptom resolve. If the condition does not improve within 7 days, seek doctor's attention.

- If you forget to use it

Apply as necessary when it is required.

- If you use too much (overdose)

In case of accidental ingestion, call Poison Control Center immediately.

While you are using it

- Things you must do

When practical, cleanse the affected area with mild soap and warm water and rinse thoroughly. Gently pat dry or air dry before application of this product. Apply externally to the affected area up to 6 times daily or after each bowel movement

- Things you must not do

Do not put this product into the rectum.

- Things to be careful of

1. Could cause burning sensation to patients with sensitivity to

Menthol.

2. Stop use and ask a doctor if:
 - i. condition worsens or does not improve within 7 days
 - ii. bleeding occurs
 - iii. symptom being treated does not subside
 - iv. redness, irritation, swelling, pain or other symptoms develop
 - v. an allergic reaction develops

Side effects

Could cause burning sensation to patients with sensitivity to Menthol.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npa.gov.my [Consumers -> Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of Calmoseptine® Ointment

- Storage

Store between 15-30°C (59-86°F)

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment

Product Description

- What it looks like

Calmoseptine® Ointment has a light pink smooth appearance with light menthol smell

- Ingredients

- Active ingredient(s)
Petrolatum, Zinc Oxide, Lanolin, Menthol

Calmoseptine[®] Ointment

Zinc Oxide(20.6%), Petrolatum(24%), Lanolin(15.7%), Menthol(0.44%)

- Inactive ingredients

Sodium borate, glycerin, phenol,
sodium bicarbonate, thymol, water,
chlorothymol, ferric oxide.

- MAL number:

MAL20046144XR

Manufacturer

Calmo Manufacturing, Inc. 16601
Burke Lane, Huntington Beach,
California 92647, USA

Product Registration Holder

Zuellig Pharma Sdn Bhd
No. 15 Persiaran Pasak Bumi, Section
U8, Perindustrian Bukit Jelutong,
40150 Shah Alam, Selangor Darul
Ehsan. Malaysia

Date of revision

23 August 2021

Serial Number

NPRA (R2) 19/224