

PERSKINDOL[®]

Classic

Gel 50 & 100 ml

Package insert

Brand or Product Name

PERSKINDOL CLASSIC GEL

Name and Strength of Active Substance(s)

Ingredients	Formula
Levomenthol	1.50% w/w
Pine Sylvestris oil	1.35% w/w
Sweet Orange oil	0.40% w/w
Wintergreen oil	0.35% w/w
Bergamot oil	0.10% w/w
Lemon oil	0.10% w/w
Terpineol	0.10% w/w
Terpinyl acetate	0.10% w/w
Rosemary oil	0.06% w/w
Lavender oil	0.02% w/w

Product Description

Perskindol Classic Gel is a white to yellowish, homogeneous gel with an aromatic odour of menthol and essential oils.

Indication

For relief of minor aches and pains of muscles and joints associated with arthritis, stiff neck, strains, sprains, bruises, cramps, simple backache.

Pharmacodynamics

Rational for combination of active ingredients:

- Menthol lowers the body temperature of the affected area therefore numbing the pain at once.
- The essential oils each have stimulant actions which help promote the blood flow to the affected area.

Recommended Dosage/Instruction

Adults and children 6 years of age and older:

Apply to affected area not more than 3 to 4 times daily.
Gently massage until PERSKINDOL is absorbed. Use only as directed.

Route of administration

For topical use.

Contraindications

Do not apply to wounds or damaged skin. Avoid contact with eyes and mucous membranes. Do not use when allergy reaction to one or more essential oils is known.

Warnings and Precautions

Keep out of reach of children. For external use only. Avoid contact with eyes and mucous membranes. Do not use with heating pads or heating devices. If condition worsens, skin redness or irritation develops or if symptoms persist for more than 7 days, discontinue use of this product and consult a doctor. Do not apply to wounds or damaged skin. Do not bandage tightly. If you have sensitive skin, consult a doctor before use. If skin irritation develops, discontinue use and consult a doctor. Do not use, pour, spill or store near heat or open flame.

Interactions

No known interactions.

Pregnancy and Lactation

As with any drug, if you are pregnant or nursing a baby, seek the advice of a health professional before using this product.

Undesirable Effects

Hypersensitivity to any of the ingredients may cause skin redness or irritation. If skin irritation develops discontinue use and consult a doctor.

Overdose and Treatment

No symptoms of overdose have been recorded.

Storage Conditions

Store below 30 °C.

Dosage forms and packaging available

100 ml tube
50 ml tube
4 ml sachet

Not all pack size available in the market.

Name and address of manufacturer

Glaropharm AG, Säntli, 8756 Miltödi, Switzerland

Product Registration Holder

Zuellig Pharma Sdn Bhd
No. 15, Persiaran Pasak Bumi Seksyen U8
Perindustrian Bukit Jelutong
40150 Shah Alam
Malaysia

MAL23016027XCRS

Date of revision

August 2023