

# COLIOPAN TABLET®

Consumer Medication Information Leaflet (RiMUP)

Butropium Bromide (5mg)

## What is in this leaflet

1. What COLIOPAN® is used for
2. How COLIOPAN® works
3. Before you use COLIOPAN®
4. How to use COLIOPAN®
5. While you are using it
6. Side effects
7. Storage and Disposal of COLIOPAN®
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

## What COLIOPAN® is used for

COLIOPAN® is indicated for spasm pains associated with inflammation or ulcer of the stomach and small intestine, formation of gallstones and any disease of the gallbladder.

## How COLIOPAN® works

COLIOPAN® relieve spasm on smooth muscle, acts to inhibit gastric secretion and has anti-ulcer effects. COLIOPAN® is also reported to act to normalize gastric emptying and to increase blood flow in stomach lining.

## Before you use COLIOPAN®

### - When you must not use it

You should not take COLIOPAN®

- if you are allergic to Butropium or any of the ingredients contained in it.
- if you have glaucoma (increased pressure within the eyeball), impaired urination due to enlargement of the prostate gland, severe heart disease, functional bowel obstruction.

### Pregnancy and lactation

Do not take COLIOPAN® if you are pregnant, trying to get pregnant or think you may be pregnant.

Do not take COLIOPAN® if you are breast-feeding. Ask your doctor or pharmacist for advice before taking any medicine.

### - Before you start to use it

Tell the doctor about all your present or past health problems. Include:

- enlargement of the prostate gland
- congestive heart failure
- irregular heartbeat
- inflammation with ulcer formation in the lining of large intestine
- overactivity of the thyroid gland

### - Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

Concurrent use of the following medicines with COLIOPAN® may intensify actions of COLIOPAN®:

- Tricyclic antidepressants
- Phenothiazines (for mental disorder)
- Monoamine oxidase (MAO) inhibitors
- Antihistamines (anti-allergy)

## How to use COLIOPAN®

### - How much to use

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

Usually, COLIOPAN® to be taken orally 2 tablets 3 times daily (30mg of Butropium bromide)

The dose should be adjusted according to your age and severity of symptoms

### - When to use it

Use as directed by your doctor or pharmacist.

### - How long to use it

Continue taking COLIOPAN® for as long as your doctor recommends.

### - If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

### - If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

## While you are using it

### - Things you must do

Tell all the doctors, dentists and pharmacists treating you that you are taking COLIOPAN®

Tell your doctor immediately if you become pregnant while taking this medication.

### - Things you must not do

Do not stop taking the medicine unless advised by your doctor. Do not take any new medicines without consulting your doctor or pharmacist.

Do not give COLIOPAN® to anyone else, even if they have the same symptoms or condition as you.

### - Things to be careful of

#### Driving and using machines

This medicine may affect your ability to drive or use machines. If the tablets make you feel sick, drowsy, dizzy or tired, or disturb your vision, do not drive or use machines and contact your doctor immediately.

## Side effects

Like all medicines, COLIOPAN® can cause side effects, although not everybody gets them.

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine.

*COLIOPAN*® may cause the following side effects:

Eyes: Symptoms of impaired eye focus may occasionally occur

Gastrointestinal: Dryness of the mouth, nausea, vomiting or constipation may occasionally occur

Genito-urinary: Impaired urination may occasionally result.

Psychoneurotic: Symptoms such as headache, drowsiness or dizziness may occasionally occur.

Cardiovascular: rapid heartbeat or low blood pressure may occasionally occur

Hypersensitivity: Hypersensitivity reactions such as skin rash may occur in rare cases, the medication should be discontinued if such signs develop

Others: Flushing or redness of the face or unusual tiredness may occasionally occur.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835490, or visiting the website [npra.gov.my](http://npra.gov.my) [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

#### **Storage and Disposal of *COLIOPAN*®**

##### **- Storage**

Keep out of the reach and sight of children.

Store below 30°C. *COLIOPAN*® tablets should be protected from moisture.

##### **- Disposal**

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

#### **Product Description**

##### **- What it looks like**

*COLIOPAN*® Tablets are hard white tablet, odorless with a bitter taste.

*COLIOPAN*® are supplied in bottles of 180 tablets

##### **- Ingredients**

- Active ingredient:  
Butropium Bromide

- Inactive ingredients:  
Lactose, Hydroxy Propyl Cellulose-L, Calcium Stearate.

##### **- MAL number:**

MAL19910111ACZ

#### **Manufacturer**

Bora Pharmaceuticals Co.,Ltd.  
No 54, Gong-Yen W. Rd  
Guan-Tyan District  
Tainan City, Taiwan

#### **Product Registration Holder**

Eisai (Malaysia) Sdn Bhd.  
Unit 701D, Level 7, Tower D, Uptown 5,  
No. 5, Jalan SS21/39, Damansara  
Uptown, 47400 Petaling Jaya,  
Selangor, Malaysia.

#### **Date of revision**

15/10/2019

#### **Serial Number**

NPRA(R4/1)031019/00285