

HIJUVEN 200 SOFT CAPSULE®

DL-Tocopheryl Nicotinate (200mg)

What is in this leaflet

1. What *Hijuven 200* is used for
2. How *Hijuven 200* works
3. Before you use *Hijuven 200*
4. How to use *Hijuven 200*
5. While you are using *Hijuven 200*
6. Side effects
7. Storage and Disposal of *Hijuven 200*
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What *Hijuven 200* is used for

Hijuven 200 is a type of Vitamin E and it is recommended for patient with the following health condition:

- High cholesterol
- Peripheral circulatory disturbances accompanying narrowing and occlusion of the arteries

How *Hijuven 200* works

DL-Tocopheryl Nicotinate is a potent Vitamin E that has shown improvement of lipid metabolism, enhancement of microcirculation, protection of vascular walls, inhibition of platelet clumping, and restoration of arterial oxygen partial pressure.

Before you use *Hijuven 200*

- When you must not use it

You must not take *Hijuven 200* if you have hypersensitivity to any substance listed in Ingredients.

Pregnancy and lactation

Do not take ***Hijuven 200*** if you are pregnant, trying to get pregnant or think you may be pregnant.

Do not take ***Hijuven 200*** if you are breast-feeding. Ask your doctor or pharmacist for advice before taking any medicine.

- Before you start to use it

Before you take *Hijuven 200*, do tell your healthcare provider if you have:

- digestive system disturbance
- dermatological problem

- Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

There is no interaction reported while using *Hijuven 200* with other medicines.

How to use *Hijuven 200*

- How much to use

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

The recommended adult dose is 3 capsules a day in three divided doses. *Hijuven 200* is to be administered orally.

Your healthcare provider may change your dose. DO NOT change your dose without talking to your healthcare provider.

- When to use it

Use as directed by your doctor or pharmacist.

- How long to use it

Dosage and duration of therapy are dependent upon individual body responses. Please stop taking *Hijuven 200* should you develop any side effects. Otherwise, *Hijuven 200* is safe for long-term consumption.

- If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the

medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

Treatment for overdose should be symptomatic and supportive.

While you are using it

- Things you must do

Take your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are taking *Hijuven 200*.

Tell your doctor immediately if you become pregnant while taking this medication.

Tell your doctor if you are lactating.

- Things you must not do

Do not stop taking the medicine unless advised by your doctor.

Do not take any new medicines without consulting your doctor.

Do not give *Hijuven 200* to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are receiving *Hijuven 200*. *Hijuven 200* may have unwanted side effects in a few people. All medicines can have side effects.

Sometimes they are serious, most of the time they are not. You may need medical attention if you get some of the side effects.

Side effects

Like all medicines, ***Hijuven 200*** can cause side effects, although not everybody gets them.

Hijuven 200 may cause the following side effects:

1. Digestive: Anorexia, gastric discomfort, stomachache, nausea, diarrhea and constipation may infrequently occur.

2. Hypersensitivity: Skin eruptions may rarely occur. Medication should be discontinued if such as symptom appears.

3. Others: Warm sensations may rarely occur.

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by ~~calling Tel: 03-78835549, or~~ visiting the website npa.gov.my ([Public → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]).

Storage and Disposal of Hijuven 200

- Storage

Keep out of the reach and sight of children.

Hijuven 200 should be stored ~~at room temperature~~ below 30°C.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

Hijuven 200 soft capsules are orange in colour. Each capsule contains 200mg of dl- α -tocopheryl nicotinate.

- Ingredients

- Active ingredient
DL-Tocopheryl Nicotinate
- Inactive ingredients

Macrogol 6000
Methylcellulose
Talc
Light Anhydrous Silicic Acid
Hydrated Silicon Dioxide
Crystalline Cellulose

- MAL number:

MAL19940373XCR

Manufacturer

Shibakawa Plant of Fuji Capsule Co., Ltd.
789 Kamiyuno, Fujinomiya-shi, Shizuoka, Japan

Product Registration Holder

Eisai (Malaysia) Sdn Bhd (18039-D)
Unit 701D, Level 7, Tower D, Uptown 5,
No. 5, Jalan SS21/39, Damansara Uptown,
47400 Petaling Jaya,
Selangor, Malaysia.

Date of revision

~~07/10/2019~~ 16/8/2022

Serial Number

~~NPRA(R2/2)220509/879NPRA(R2/1)191007/263~~