

# BEOVA™ 50mg Film Coated Tablets

Vibegron 50mg

Consumer Medication Information Leaflet  
(RiMUP)

## What is in this leaflet

1. What Beova is used for
2. How Beova works
3. Before you use Beova
4. How to use Beova
5. While you are using it
6. Side effects
7. Storage and Disposal of Beova
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision
11. Serial number

## What Beova is used for

Beova is a prescription medicine for adults use to treat the following symptoms due to a condition called overactive bladder:

- urge urinary incontinence: a strong need to urinate with leaking or wetting accidents
- urgency: the need to urinate right away
- frequency: urinating often

## How Beova works

Beova selectively stimulates the relaxation of muscles in the bladder to promote urine accumulation function, and improves urinary urgency, urinary frequency and urge incontinence in overactive bladder patients.

## Before you use Beova

### - When you must not use it

If you are allergic to vibegron or any of the ingredients of this medicine.

### *Pregnancy and lactation*

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. It is not known if Beova will harm your unborn baby or passes into your breast milk.

It is also not known if Beova is safe and effective in children as there is no clinical studies have been conducted.

### - Before you start use it

Consult your doctor before using Beova:

- If you have liver problems
- If you have kidney problems
- If you have severe heart disease

- If you have trouble emptying your bladder

### - Taking other medicines

Tell your doctor or pharmacist if you are taking or have recently taken or might take any other medicines. Ask your doctor for advice before taking any of the following medicines together with Beova:

- Antifungal medicine such as itraconazole, ketoconazole
- Medicines for HIV infection such as Ritonavir
- Antibacterial medicine such as rifampicin
- Antiseizure medicine such as phenytoin, carbamazepine
- Medicines that contain digoxin

## How to use Beova

### - How much to use

Always take this medicine exactly as your doctor has told you. Check with your doctor if you are not sure.

The usual dosage for oral use is 50 mg once daily after meal.

### - When to use it

Use only as directed by your physician.

### - How long to use it

Continue taking Beova for as long as your doctor recommends.

### - If you forget to use it

Do not take a double dose to make up for a forgotten dose. Skip the missed dose and take the next dose at the time you normally take it.

### - If you use too much (overdose)

Contact your doctor immediately and take the Beova pack with you.

## While you are using it

### - Things you must do

Do not stop taking Beova without first talking to your doctor.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

### - Things you must not do

Do not take any new medicines without consulting your doctor.

## Side effects

Though incidence is unknown, Beova may cause significant side effect like inability to empty your bladder (urinary retention). It may increase your chances of not being able to empty your bladder, especially if you have bladder outlet obstruction or take other medicines for treatment of overactive bladder. Tell your doctor right away if you are unable to empty your bladder.

The following side effects also may occur. Therefore, patients should be closely monitored and if any abnormal findings are observed, consult the doctor for appropriate measures.

Common:

- Dry mouth, constipation, urinary tract infection, increased urine retained in the bladder.

Uncommon:

- Eye discomfort in bright light, increased liver enzyme.

Frequency unknown:

- Headache, dizziness, trouble sleeping, drowsiness, nausea, diarrhea, abdominal pain, decreased appetite, gastric, abnormal heart rhythm, palpitation, difficulty in urination, bladder pain, bedwetting, pain during urination, rash, excessive sweating, skin itching, dry eye, blurred vision, abnormal liver function, fatigue, hot flush, high cholesterol, edema, muscle aches, thirst, increased blood pressure, chest discomfort.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website [npa.gov.my](http://npa.gov.my) [Consumers→ Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

## Storage and Disposal of Beova

### - Storage

Do not store above 30°C

### - Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of

# BEOVA™ 50mg Film Coated Tablets

Consumer Medication Information Leaflet  
(RiMUP)

Vibegron 50mg

---

medicines no longer required. These measures will help to protect the environment.

## Product Description

### - What it looks like

Beova is a round light green film-coated tablet with “BV50” on one side.

### - Ingredients

- Active ingredient
  - Vibegron
- Excipients
  - D-Mannitol
  - Microcrystalline cellulose
  - Croscarmellose sodium
  - Hydroxypropyl cellulose
  - Magnesium stearate
  - Lactose hydrate
  - Hypromellose
  - Titanium oxide
  - FD&C Blue No.2 Aluminium Lake
  - Triacetin
  - Yellow ferric oxide
  - Carnauba wax

### - MAL number:

## Manufacturer

KYORIN Pharmaceutical Group  
Facilities Co., Ltd. Noshiro Plant,  
1, Matsubara, Noshiro-shi,  
Akita 016-0000, Japan

## Packager

Bora Pharmaceuticals Co., Ltd.,  
No.54, Gong-Yeh W. Rd., Guan-Tyan  
District, Tainan City, Taiwan

## Product Registration Holder

Eisai (Malaysia) Sdn. Bhd.,  
Unit 701D, Level 7, Tower D, Uptown 5,  
No. 5, Jalan SS21/39, Damansara  
Uptown, 47400 Petaling Jaya,  
Selangor, Malaysia

## Date of revision

28/10/2024

## Serial number

NPRA (R1) 24/005