

This insert is applicable to all pack size



## NCHK BABAO BAO YING DAN

**Indication:** Traditionally used for reducing crying at night, fever, cold & cough.  
**Dosage:** Children 2-3 years old: 1 small spoonful.  
Children 4-10 years old: 2 small spoonful.  
To be taken 2 times per day before or after meal with warm water

**Indikasi:** Digunakan secara tradisional untuk mengurangkan menangis pada malam, demam, selsema & batuk.

**Dos:** Kanak – kanak 2-3 tahun, 1 sudu kecil.  
Kanak-kanak 4-10 tahun, 2 sudu kecil teh.  
Makan 2 kali sehari sebelum atau selepas makan dengan air suam.

### Each 1g of powder contains:

川贝母 Bulbus Fritillariae Cirrhosae	200mg
珍珠母 Concha Margaritifera Usta	70mg
牛黄 Calculus Bovis	70mg
黄芩 Radix Scutellaria Baicalensis	150mg
茯苓 Sclerotium Poria Cocos Wolf	160mg
羌活 Rhizoma Notopterygium Incisum	200mg
天竺黄 Concretio Silicae Bambusae	80mg
桑叶 Folium Morus Alba	70mg

This is a traditional medicine. Please consult your pharmacist/doctor if symptoms persist or worsen. Keep out of reach of children. Jauhkan daripada capaian kanak-kanak. Store below 30°C, protect from light and moisture. Pregnancy and breastfeeding: Insufficient reliable data. This product contains animal part. This product contains substances from animal origin.



## 八寶 保嬰丹



NCHK046

主治：传统上用于减少夜啼，发烧，感冒及咳嗽。

服法：小孩2-3岁：1茶匙。  
小孩4-10岁：2茶匙。  
每日两次，饭前饭后用温水送服。

**Deskripsi Produk:** Serbuk halus yang berwarna perang. Tidak berbau dan berasa sedikit pahit.

Manufactured by:  
**TJ-TYT PHARMACEUTICALS (M) SDN. BHD.**  
1 & 3, Persiaran Perindustrian Pengkalan 12,  
Kawasan Perindustrian Pengkalan,  
31500 Lahat, Perak, Malaysia.

Distributed by Product Registration Holder:  
**NCHK Medicine Enterprise Sdn. Bhd.**  
21G, Jalan SL 11/7, Bandar Sg. Long,  
43000 Kajang, Selangor.

Available Pack Size: 3gm / 5gm

Revision Date: 18 Dec 2024



046MY124-02V