

Directions for Use

Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion

1. NAME OF THE MEDICINAL PRODUCT

Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each ml of solution for injection / infusion contains 10 mg rocuronium bromide. Each ampoule with 5 ml contains 50 mg rocuronium bromide. For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Solution for injection / infusion

Product description:

Clear, colourless to pale brownish-yellow solution
pH of the solution: 3.8 to 4.2
Osmolality: 270 – 310 mOsmol/kg.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion is indicated as an adjunct to general anaesthesia to facilitate endotracheal intubation, to provide skeletal muscle relaxation and to facilitate mechanical ventilation in adults, children and infants from one month of age. Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion is also indicated as an adjunct in the intensive care unit (ICU) to facilitate mechanical ventilation as part of Rapid Sequence Induction, however, this has not been studied in infants and children. See also section 4.2 and 5.1.

4.2 Posology and method of administration

Posology

Like other neuromuscular blocking agents, Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion should only be administered by, or under supervision of, experienced clinicians who are familiar with the action and use of these drugs.

As with other neuromuscular blocking agents, the dosage of Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion should be individualised in each patient. The method of anaesthesia and the expected duration of surgery, the method of sedation and the expected duration of mechanical ventilation, the possible interaction with other medicinal products that are administered concomitantly and the condition of the patient should be taken into account when determining the dose. The use of an appropriate neuromuscular monitoring technique is recommended for the evaluation of the neuromuscular block and recovery.

Surgical procedures

Inhalational anaesthetics potentiate the neuromuscular blocking effects of Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion. This potentiation becomes clinically relevant during the course of anaesthesia when a certain tissue concentration of the volatile agents is reached. Consequently, adjustments should be made by administering smaller maintenance doses at less frequent intervals or by using lower infusion rates of Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion during long lasting procedures (longer than 1 hour) under inhalational anaesthesia (see 4.5).

Adults

In adult patients the following dosage recommendations may serve as a general guidance for tracheal intubation and muscle relaxation for short to long lasting surgical procedures and for use in the intensive care unit. This medicinal product is for single use only.

Tracheal intubation:

The standard intubating dose during routine anaesthesia is 0.6 mg Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion per kg body weight, which results in adequate intubation conditions within 60 seconds in nearly all patients. A dose of 1.0 mg Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion per kg body weight is recommended for facilitating tracheal intubation conditions during rapid sequence induction of anaesthesia, after which adequate intubation conditions are also established within 60 seconds in nearly all patients. If a dose of 0.6 mg Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion per kg body weight is used for rapid sequence induction of anaesthesia, it is recommended to intubate the patient 90 seconds after administration of Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion

Maintenance dosage:

The recommended maintenance dose is 0.15 mg Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion per kg body weight. In case of long-term inhalational anaesthesia it should be reduced to 0.075 – 0.1 mg of Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion per kg body weight. The maintenance doses should best be given when twitch height has recovered to 25 % of control twitch height, or when 2 to 3 responses to train-of-four stimulation (TOF) are present.

Continuous infusion:

If Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion is administered by continuous infusion, it is recommended to give a loading dose of 0.6 mg Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion per kg body weight and, when the neuromuscular block starts to recover, to start administration by infusion. The infusion rate should be adjusted to maintain twitch response at 10 % of control twitch height or to maintain 1 to 2 responses to train-of-four stimulation.

In adults under intravenous anaesthesia, the infusion rate required to maintain the neuromuscular block at this level ranges from 0.3 – 0.6 mg/kg/h. Under inhalational anaesthesia the infusion rate ranges from 0.3 – 0.4 mg/kg/h. Continuous monitoring of the neuromuscular block is essential since infusion rate requirements vary from patient to patient and with the anaesthetic method used.

Dosage in pregnant patients:

In patients undergoing Caesarean section, it is recommended to only use a dose of 0.6 mg Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion per kg body weight, since a 1.0 mg/kg dose has not been investigated in this patient group. Reversal of neuromuscular block induced by neuromuscular blocking agents may be inhibited or unsatisfactory in patients receiving magnesium salts for toxemia of pregnancy because magnesium salts enhance neuromuscular blockade. Therefore, in these patients the dosage of rocuronium should be reduced and be titrated to twitch response.

Paediatric population:

For infants (28 days – 23 months), children (2 – 11 years) and adolescents (12 – 18 years) the recommended intubation dose during routine anaesthesia and maintenance dose are similar to those in adults.

For continuous infusion in paediatrics, the infusion rates, with exception of children, are the same as for adults. For children higher infusion rates might be necessary. Thus, for children the same initial infusion rates as for adults are recommended and then this should be adjusted to maintain twitch response at 10 % of control twitch height or to maintain 1 or 2 responses to train of four stimulation during the procedure. There are insufficient data to support dose recommendations for the use of Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion in neonates (0 – 1 month). The experience with Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion in rapid sequence induction in paediatric patients is limited. Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion is therefore not recommended for facilitating tracheal intubation conditions during rapid sequence induction in paediatric patients.

Elderly patients and patients with hepatic and/or biliary tract disease and/or renal failure: The standard intubation dose for geriatric patients and patients with hepatic and/or biliary tract disease and/or renal failure during routine anaesthesia is 0.6 mg Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion per kg body weight. A dose of 0.6 mg per kg body weight should be considered for rapid sequence induction of anaesthesia in patients in which a prolonged duration of action is expected however adequate conditions for intubation may not be established for 90 seconds after administration of Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion. Regardless of the anaesthetic technique used, the recommended maintenance dose for these patients is 0.075 – 0.1 mg Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion per kg body weight, and the recommended infusion rate is 0.3 – 0.4 mg/kg/h (see also Continuous infusion) (see also section 4.4).

Overweight and obese patients:

When used in overweight or obese patients (defined as patients with a body weight of 30 % or more above ideal body weight) doses should be reduced taking into account the lean body mass.

Intensive care procedures

Tracheal intubation

For tracheal intubation, the same doses should be used as described above under surgical procedures.

Maintenance dosing

The use of an initial loading dose of 0.6 mg·kg⁻¹ Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion is recommended, followed by a continuous infusion as soon as twitch height recovers to 10 % or upon reappearance of 1 to 2 twitches to train of four stimulation. Dosage should always be titrated to effect in the individual patient. The recommended initial infusion rate for the maintenance of a neuromuscular block of 80 – 90 % (1 to 2 twitches to TOF stimulation) in adult patients is 0.3 – 0.6 mg·kg⁻¹·h⁻¹ during the first hour of administration, which will need to be decreased during the following 6 – 12 hours, according to the individual response. Thereafter, individual dose requirements remain relatively constant.

A large between patient variability in hourly infusion rates has been found in controlled clinical studies, with mean hourly infusion rates ranging from 0.2 – 0.5 mg·kg⁻¹·h⁻¹ depending on nature and extent of organ failure(s), concomitant medication and individual patient characteristics. To provide optimal individual patient control, monitoring of neuromuscular transmission is strongly recommended. Administration up to 7 days has been investigated.

Special populations

Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion is not recommended for the facilitation of mechanical ventilation in the intensive care in pediatric and geriatric patients due to a lack of data on safety and efficacy.

Method of administration

Intravenous use

Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion is administered intravenously (i.v.) either as a bolus injection or as a continuous infusion (for further information see also section 6.6).

Risk of medication errors

Accidental administration of neuromuscular blocking agents may result in serious adverse events, including fatal outcomes

4.3 Contraindications

Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion is contra-indicated in patients with hypersensitivity to rocuronium or to the bromide ion or to any of the excipients listed in section 6.1.

4.4 Special warnings and precautions for use

Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion should be administered only by experienced staff familiar with the use of neuromuscular blocking agents. Adequate facilities and staff for endotracheal intubation and artificial ventilation have to be available for immediate use.

Since Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion causes paralysis of the respiratory muscles, ventilatory support is mandatory for patients treated with this active substance until adequate spontaneous respiration is restored. As with all neuromuscular blocking agents, it is important to anticipate intubation difficulties, particularly when used as part of a rapid sequence induction technique.

As with other neuromuscular blocking agents, residual curarization has been reported for Rocuronium. In order to prevent complications resulting from residual curarization, it is recommended to extubate only after the patient has recovered sufficiently from neuromuscular block. Other factors which could cause residual curarization after extubation in the post-operative phase (such as drug interactions or patient condition) should also be considered. If not used as part of standard clinical practice, the use of a reversal agent (such as sugammadex or acetylcholinesterase inhibitors) should be considered, especially in those cases where residual curarization is more likely to occur. It is essential to ensure that the patient is breathing spontaneously, deeply and regularly before leaving the theatre after anaesthesia.

Anaphylactic reactions (see above) can occur after the administration of neuromuscular blocking agents. Precautions for treating such reactions should always be taken. Particularly in the case of previous anaphylactic reactions to neuromuscular blocking agents, special precautions should be taken since allergic cross-reactivity to neuromuscular blocking agents has been reported.

Dose levels higher than 0.9 mg Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion per kg body weight may increase the heart rate; this effect could counteract the bradycardia produced by other anaesthetic agents or by vagal stimulation.

In general, following long term use of muscle relaxants in the ICU, prolonged paralysis and/or skeletal muscle weakness has been noted. In order to help preclude possible prolongation of neuromuscular blockage and/or overdose, it is strongly recommended that neuromuscular transmission is monitored throughout the use of muscle relaxants. In addition, patients should receive adequate analgesia and sedation. Furthermore, muscle relaxants should be titrated to the effect in the individual patient. This should be done by or under the supervision of experienced clinicians who are familiar with the effects and with appropriate neuromuscular monitoring techniques.

Myopathy has been regularly reported after long-term concurrent use of non-depolarising neuromuscular blockers and corticosteroids. The co-administration period should be reduced to be as short as possible (see section 4.5).

If suxamethonium is used for intubation, the administration of rocuronium should be delayed until the patient has clinically recovered from the neuromuscular block induced by suxamethonium.

The following conditions may influence the pharmacokinetics and/or pharmacodynamics of Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion:

Hepatic and/or biliary tract disease and renal failure

Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion is excreted in urine and bile. Therefore, it should be used with caution in patients with clinically significant hepatic and/or biliary diseases and/or renal failure. In these patient groups prolongation of the effect has been observed with doses of 0.6 mg Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion per kg body weight.

Prolonged circulation time

Conditions associated with prolonged circulation time such as cardiovascular diseases, advanced age and an oedematous state resulting in an increased volume of distribution, may contribute to a slower onset of the effect. The duration of action may also be prolonged due to a reduced plasma clearance.

Neuromuscular disease

Like other neuromuscular blocking agents, Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion should be used with extreme caution in patients with a neuromuscular disease or after poliomyelitis since the response to neuromuscular blocking agents may be considerably altered in these cases. The magnitude and direction of this alteration may vary widely. In patients with myasthenia gravis or with the myasthenic (Eaton-Lambert) syndrome, small doses of Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion may have profound effects and Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion should be titrated to the response.

Hypothermia

In surgery under hypothermic conditions, the neuromuscular blocking effect of Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion is increased and the duration prolonged.

Obesity

Like other neuromuscular blocking agents, Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion may exhibit a prolonged duration and a prolonged spontaneous recovery in obese patients, when the administered doses are calculated on actual body weight.

Burns

Patients with burns are known to develop resistance to non-depolarising neuromuscular blocking agents. It is recommended that the dose is titrated to the response.

Conditions which may increase the effects of Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion

Hypokalaemia (e.g. after severe vomiting, diarrhoea or diuretic therapy), hypermagnesaemia, hypocalcaemia (after massive transfusions), hypoproteinaemia, dehydration, acidosis, hypercapnia and cachexia. Severe electrolyte disturbances, altered blood pH or dehydration should therefore be corrected when possible.

This medicinal product contains less than 1 mmol sodium (23 mg) per dose, i.e. essentially 'sodium-free'.

4.5 Interaction with other medicinal products and other forms of interaction

The following medicinal products have been shown to influence the magnitude and/or duration of the effect of non-depolarising neuromuscular blocking agents:

Effect of other medicinal products on Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion

Increased effect:

- Halogenated volatile anaesthetics potentiate the neuromuscular block for Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion. The effect only becomes apparent with maintenance dosing (see section 4.2). Reversal of the block with acetylcholinesterase inhibitors could also be inhibited
- High doses of: thiopental, methohexital, ketamine, fentanyl, gammahydroxybutyrate, etomidate and propofol
- After intubation with suxamethonium. (see section 4.4).
- Long term concomitant use of corticosteroids and Rocuronium in the ICU may result in prolonged duration of neuromuscular block or myopathy (see sections 4.4 and 4.8).

Other medicinal products:

- antibiotics: aminoglycosides, lincosamides (e.g. lincomycin and clindamycin), polypeptide antibiotics, acylamino-penicillin antibiotics, tetracyclines, high doses metronidazole.
- diuretics, quinidine and its isomer quinine, protamine, adrenergic blocking agents, magnesium salts, calcium channel blocking agents and lithium salts and local anaesthetics (lidocaine i.v., bupivacaine epidural) and acute administration of phenytoin or β -blocking agents.

Recurarization has been reported after post-operative administration of: aminoglycoside, lincosamide, polypeptide and acylamino-penicillin antibiotics, quinidine, quinine and magnesium salts (see section 4.4).

Decreased effect:

- Neostigmine, edrophonium, pyridostigmine, aminopyridine derivatives
- Prior chronic administration of phenytoin or carbamazepine
- protease inhibitors

Variable effect:

- Administration of other non-depolarising neuromuscular blocking agents in combination with Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion may produce attenuation or potentiation of the neuromuscular block, depending on the order of administration and the neuromuscular blocking agent used.
- Suxamethonium given after the administration of Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion may produce potentiation or attenuation of the neuromuscular blocking effect of Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion.

Effect of rocuronium on other drugs:

Combined use with lidocaine could result in a faster onset of the effect of lidocaine.

Paediatric population

No formal interaction studies have been performed. The above mentioned interactions for adults and their special warnings and precautions for use (see section 4.4) should also be taken into account for paediatric patients.

4.6 Fertility, pregnancy and lactation

Pregnancy

There are limited data on the use of Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion in pregnant women. Animal studies do not indicate direct or indirect harmful effects with respect to reproductive toxicity (see section 5.3). Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion should only be given to pregnant women when strictly necessary and the attending physician decides that the benefits outweigh the risks.

Cesarean section

In patients undergoing Caesarean section, Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion can be used as part of a rapid sequence induction technique, provided no intubation difficulties are anticipated and a sufficient dose of anaesthetic agent is administered or following suxamethonium facilitated intubation. Use of Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion during caesarean section has been shown to be safe at doses of 0.6 mg/kg bodyweight. It does not affect the Apgar score, the foetal muscle tone or the cardiorespiratory adaptation. From umbilical cord blood sampling it is apparent that only limited placental transfer of Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion occurs, which does not lead to the observation of clinical adverse reactions in the new-born infant.

Note 1: doses of 1.0 mg/kg have been investigated during rapid sequence induction of anaesthesia, but not in Caesarean section patients.

Note 2: reversal of neuromuscular block induced by neuromuscular blocking agents may be inhibited or unsatisfactory in patients receiving magnesium salts for toxemia of pregnancy because magnesium salts enhance neuromuscular blockade. Therefore, in these patients the dosage of Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion should be reduced and be titrated to twitch response.

Breast-feeding

It is unknown whether Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion / metabolites are excreted in human milk. Other medicinal products of this class show little excretion into breast milk and low resorption by the suckling child. Animal studies have shown excretion of Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion in insignificant amounts in breast milk.

A decision must be made whether to discontinue breast-feeding or to discontinue/abstain from Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion therapy taking into account the benefit of breast-feeding for the child and the benefit of therapy for the woman.

Fertility

There are no data with regard to effects of Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion on fertility.

4.7 Effects on ability to drive and use machines

Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion has major influence on the ability to drive and use machines. It is not recommended to use potentially dangerous machinery or to drive a car during the first 24 hours after the full recovery from the neuromuscular blocking action of Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion. Since Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion is used as an adjunct to general anaesthesia, the usual precautionary measures after a general anaesthesia should be taken for ambulatory patients.

4.8 Undesirable effects

The frequency of undesirable effects is classified into the following categories:

Uncommon/rare	($>1/10\,000$ to $<1/100$)
Very rare	($<1/10,000$)
Not known	(frequency cannot be estimated from the available data)

Frequencies are estimates derived from post-marketing surveillance reports and data from the general literature. Post-marketing surveillance data cannot give precise incidence figures. For that reason, the reporting frequency was divided over three rather than five categories.

The most common undesirable effects are pain and/or local reactions around injection site, changes in vital functions and prolonged neuromuscular block. The most frequently reported serious adverse drug reactions during post-marketing surveillance is 'anaphylactic and anaphylactoid reactions' and associated symptoms. See also the explanations below.

Approval for Printing

B | BRAUN Melsungen AG

Approved for Printing

Approved for Printing when corrected

New draft required

LRA:

Date _____ Signature _____

Name in capital letters _____

LMS:

Date _____ Signature _____

Name in capital letters _____

Black

Dimension = 210 x 594 mm

2 pages

LLD-Spec.: L94

Lätus 8468



MY___0709

0709/12628396/1024

LF (MPc)

Production site: Berlin

Font size: 8 pt.

G 201038



Immune system disorders*Very rare:*

- Hypersensitivity
- Anaphylactic reaction *
- Anaphylactoid reaction*
- Anaphylactic shock
- Anaphylactoid shock

Nervous system disorders*Very rare:*

- Flaccid paralysis

Eye disorders*Not known*

- Mydriasis, fixed pupils (in the context of a potential increase of permeability or compromise of the integrity of the Blood Brain Barrier (BBB))

Cardiac disorders*Uncommon/rare:*

- Tachycardia

Not known:

- Kounis syndrome

Vascular disorders*Uncommon/rare:*

- Hypotension

Very rare:

- Circulatory collapse and shock
- Flushing

Respiratory, thoracic and mediastinal disorders*Very rare:*

- Bronchospasm

Not known

- Apnoea
- Respiratory failure

Skin and subcutaneous tissue disorders*Very rare:*

- Rash, erythematous rash
- Angioneurotic oedema
- Urticaria
- Itching
- Exanthema

Musculoskeletal, connective tissue and bone disorders*Very rare:*

- skeletal muscle weakness (after long-term use in the ICU)
- steroid myopathy (after long-term use in the ICU) (see section 4.4)

General disorders and administration site conditions*Very rare:*

- Face oedema

Uncommon/rare:

- Drug ineffective
- Drug effect/ therapeutic response decreased
- Drug effect/ therapeutic response increased
- Injection site pain and/or local reactions*

Investigations*Very rare:*

- Increased histamine level*

Injury, poisoning and procedural complications*Very rare:*

- Airway complication of anaesthesia

Uncommon/rare:

- Prolonged neuromuscular block*
- Delayed recovery from anaesthesia

Paediatric population

A meta-analysis of 11 clinical studies in paediatric patients (n = 704) with Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion (up to 1 mg/kg) showed that tachycardia was identified as adverse drug reaction with a frequency of 1.4 %.

Information on particular undesirable effects*Anaphylaxis**

Although very rare, severe anaphylactic/anaphylactoid reactions to neuromuscular blocking agents including Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion have been reported. Anaphylactic/anaphylactoid reactions are: bronchospasm, cardiovascular changes (e.g. hypotension, tachycardia, circulatory collapse – shock), and cutaneous changes (e.g. angioedema, urticaria). These reactions have, in some cases been fatal. Due to the possible severity of these reactions, one should always assume they may occur and take the necessary precautions.

Local injection site reactions

During rapid sequence induction of anaesthesia, pain on injection has been reported, especially when the patient has not yet completely lost consciousness and particularly when propofol is used as the induction agent. In clinical studies, pain on injection has been noted in 16 % of the patients who underwent rapid sequence induction of anaesthesia with propofol and in less than 0.5 % of the patients who underwent rapid sequence induction of anaesthesia with fentanyl and thiopental.

Increased histamine level

Since neuromuscular blocking agents are known to be capable of inducing histamine release both locally at the site of injection and systemically, the possible occurrence of itching and erythematous reaction at the site of injection and/or generalised histaminoid (anaphylactoid) reactions (see also under anaphylactic reactions above) should always be taken into consideration when administering these drugs. In clinical studies only a slight increase in mean plasma histamine level has been observed following rapid bolus administration of 0.3 – 0.9 mg Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion per kg body weight.

Prolonged neuromuscular block

The most frequent adverse reaction to non-depolarising blocking agents as a class consists of an extension of the agent's pharmacological action beyond the time period needed. This may vary from skeletal muscle weakness to profound and prolonged skeletal muscle paralysis resulting in respiratory insufficiency or apnoea.

Myopathy

Myopathy has been reported after the use of various neuromuscular blocking agents in the ICU in combination with corticosteroids (see section 4.4).

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in Appendix V.

4.9 Overdose**Symptoms**

The main symptom of overdose is prolonged neuromuscular block.

In animal studies, severe depression of cardiovascular function, ultimately leading to cardiac collapse, did not occur until a cumulative dose of 750 x ED₅₀ (135 mg Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion per kg body weight) was administered.

Treatment

In the event of overdose and prolonged neuromuscular block, the patient should continue to receive ventilatory support and sedation. There are two options for the reversal of neuromuscular block: (1) In adults, sugammadex can be used for reversal of intense (profound) and deep block. The dose of sugammadex to be administered depends on the level of neuromuscular block. (2) An acetylcholinesterase inhibitor (e.g. neostigmine, edrophonium, pyridostigmine) or sugammadex can be used once spontaneous recovery starts and should be administered in adequate doses. When administration of an acetylcholinesterase inhibiting agent fails to reverse the neuromuscular effects of Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion, ventilation must be continued until spontaneous breathing is restored. Repeated dosages of an acetylcholinesterase inhibitor can be dangerous.

5. PHARMACOLOGICAL PROPERTIES**5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: muscle relaxants, peripherally acting agents, other quaternary ammonium compounds.
ATC code: M03AC09

Pharmacodynamic effects

Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion is an intermediate acting, non-depolarising neuromuscular blocking agent with a fast onset, possessing all of the characteristic pharmacological actions of this class of medicinal products (curariform). It acts by competing for nicotinic cholinceptors at the motor end-plate. The ED₅₀ (dose required to produce 90 % depression of the twitch response of the thumb to stimulation of the ulnar nerve) during balanced anaesthesia is approximately 0.3 mg per kg body weight.

Clinical efficacy and safety**Routine practice**

Within 60 seconds after intravenous administration of a dose of 0.6 mg Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion per kg body weight (2 x ED₅₀ under balanced anaesthesia), adequate intubation conditions can be achieved in nearly all patients. In 80 % of these patients intubation conditions are rated excellent. Within 2 min general muscle paralysis adequate for any type of procedure is established. The clinical duration (the duration until spontaneous recovery to 25 % of control twitch height with this dose is 30 – 40 min. The total duration (time until spontaneous recovery to 90 % of control twitch height) is 50 min. The mean time of spontaneous recovery of twitch response from 25 to 75 % (recovery index) after a bolus dose of 0.6 mg Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion per kg body weight is 14 min.

With lower dosages of 0.3 – 0.45 mg Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion per kg body weight (1 – 1 ½ x ED₅₀), the onset of the effect is slower and the duration of action is shorter (13 – 26 min). After administration of 0.45 mg Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion per kg body weight, acceptable intubation conditions are reached after 90 seconds.

Emergency intubation

During rapid sequence induction of anaesthesia under propofol or fentanyl/thiopental anaesthesia, adequate intubation conditions are achieved within 60 seconds in 93 % and 96 % of the patients respectively, after administration of a dose of 1.0 mg Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion per kg body weight. Of these, 70 % are rated excellent. The clinical duration with this dose approaches 1 hour, at which time the neuromuscular block can be safely reversed. After administration of a dose of 0.6 mg Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion per kg body weight, adequate intubation conditions are achieved within 60 seconds in 81 % and 75 % of the patients during a rapid sequence induction technique with propofol or fentanyl/thiopental, respectively.

Doses higher than 1.0 mg Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion per kg body weight will not improve the intubation conditions appreciably; the duration of the effect, however, will be prolonged. Doses higher than 4 x ED₉₀ have not been studied.

Intensive care

The use of rocuronium in the Intensive Care Unit was studied in two open-label trials. A total of 95 adult patients were treated with an initial dose of 0.6 mg Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion per kg body weight, followed by a continuous infusion of 0.2 – 0.5 mg/kg/h during the first hour of administration as soon as twitch height recovers to 10 % or upon reappearance of 1 to 2 twitches to train-of-four (TOF) stimulation. The dosages were individually titrated. In the following hours, doses were decreased under regular monitoring of the TOF stimulation. Administration for a time period of up to 7 days has been investigated.

Adequate neuromuscular blockade was achieved, but a high variability in hourly infusion rates between patients and a prolonged recovery from neuromuscular blockade was observed.

The time to recover of the train of four ratio to 0.7 is not significantly correlated to the total duration of rocuronium infusion. After a continuous infusion for 20 hours or more the median (range) time between return of T2 to train of four stimulation and recovery of the train of four ratio to 0.7 varied between 0.8 and 12.5 hours in patients without multiple organ failure and 1.2 – 25.5 hours in patients with multiple organ failure.

Paediatric population

Mean onset time in infants, toddlers and children at an intubation dose of 0.6 mg/kg is slightly shorter than in adults. Comparison within paediatric age groups showed that the mean onset time in neonates and adolescents (1 min) is slightly longer than in infants, toddlers and children (0.4, 0.6 and 0.8 min, respectively).

The duration of relaxation and the time to recovery tend to be shorter in children compared to infants and adults. Comparing within paediatric age groups demonstrated that mean time to reappearance of T3 was prolonged in neonates and infants (56.7 and 60.7 min, respectively) when compared to toddlers, children and adolescents (45.3, 37.6 and 42.9 min, respectively).

Mean (SD) time to onset and clinical duration following 0.6 mg/kg rocuronium initial intubating dose during sevoflurane/nitrous oxide and isoflurane/nitrous oxide (maintenance) anaesthesia (Paediatric patients) PP group*

	Time to maximum block ** (min)	Time to reappearance of T3 ** (min)
Neonates (0 – 27 days) n = 10	0.98 (0.62)	56.69 (37.04) n = 9
Infants (28 days – 2 months) n = 11	0.44 (0.19) n = 10	60.71 (16.52) n = 11
Toddler (3 months – 23 months) n = 30	0.59 (0.27) n = 28	45.46 (12.94) n = 27
Children (2 – 11 years) n = 34	0.84 (0.29) n = 34	37.58 (11.82)
Adolescents (12 – 17 years) n = 31	0.98 (0.38)	42.90 (15.83) n = 30

* Dose of rocuronium administered within 5 seconds.

** Calculated from the end of administration of the rocuronium intubating dose

Special populations

The duration of the effect of maintenance doses of 0.15 mg Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion per kg body weight might be somewhat longer under enflurane and isoflurane anaesthesia in geriatric patients and in patients with hepatic or renal disease (approximately 20 min) than in patients without impairment of excretory organ functions under intravenous anaesthesia (approximately 13 min). No cumulation of effect (progressive increase in duration of action) with repetitive maintenance doses at the recommended level has been observed.

Cardiovascular surgery

In patients scheduled for cardiovascular surgery the most common cardiovascular changes during the onset of maximum blockage after receiving a dose of 0.6 – 0.9 mg Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion per kg body weight are a slight and clinically insignificant increase in heart rate up to 9 % and an increase in mean arterial blood pressure up to 16 % from the control values

Antagonists

Administration of acetylcholinesterase inhibitors, such as neostigmine, pyridostigmine or edrophonium, antagonises the action of Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion.

5.2 Pharmacokinetic properties**Distribution and elimination**

After intravenous administration of a single bolus dose of Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion, the time course of the plasma concentration runs in three exponential phases. In normal adults, the mean (95 % CI) elimination half-life is 73 (66 – 80) min, the (apparent) volume of distribution at steady state conditions is 203 (193 v 214) ml/kg and the plasma clearance is 3.7 (3.5 – 3.9) ml/kg/min.

When administered as a continuous infusion to facilitate mechanical ventilation for a time period of 20 hours or more, the mean elimination half-life and the mean (apparent) volume of distribution at steady state are increased. A high variability between patients was found in controlled clinical studies, related to the nature and extent of (multiple) organ failure and individual patient characteristics. In patients with multiple organ failure a mean (± SD) elimination half-life of 21.5 (± 3.3) hours, an (apparent) volume of distribution at steady state of 1.5 (± 0.8) l·kg⁻¹ and a plasma clearance of 2.1 (± 0.8) ml/kg/min were found.

Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion is excreted in urine and bile. Excretion in urine approaches 40 % within 12 – 24 hours. After injection of a radiolabeled dose of Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion, excretion of the radiolabel is on average 47 % in urine and 43 % in faeces after 9 days. Approximately 50 % is recovered as Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion.

Biotransformation

No metabolites are detected in the plasma.

Paediatric population

The apparent volume of distribution in infants (3 – 12 months) is higher compared to older children (1 – 8 years) and adults. In children aged 3 – 8 years, clearance is higher and the elimination half-life is approximately 20 min shorter compared to adults and children < 3 years.

Pharmacokinetics (PK) of Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion in paediatric patients (n = 146) with ages ranging from 0 to 17 years were evaluated using a population analysis of the pooled pharmacokinetic datasets from two clinical trials under sevoflurane (induction) and isoflurane/nitrous oxide (maintenance) anaesthesia. All pharmacokinetic parameters were found to be linearly proportional to body weight illustrated by a similar clearance (CL; l/kg/h). The volume of distribution (l/kg) and elimination half-life (h) decrease with age (years). The pharmacokinetic parameters of typical paediatrics within each age group are summarized below:

Estimated PK parameters of Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion in typical paediatric patients during sevoflurane and nitrous oxide (induction) and isoflurane/Nitrous oxide (maintenance anaesthesia)

PK Parameters	Patient Age Range				
	Term newborn infants (0 – 27 days)	Infants (28 days to 2 months)	Toddlers (3 – 23 months)	Children (2 – 11 years)	Adolescents (12 – 17 years)
CL (l/kg/h)	0.31 (0.07)	0.30 (0.08)	0.33 (0.10)	0.35 (0.09)	0.29 (0.14)
Volume of Distribution (l/kg)	0.42 (0.06)	0.31 (0.03)	0.23 (0.03)	0.18 (0.02)	0.18 (0.01)
t _{1/2β} (h)	1.1 (0.2)	0.9 (0.3)	0.8 (0.2)	0.7 (0.2)	0.8 (0.3)

Geriatric patients and patients with renal dysfunction

The plasma clearance in geriatric patients and in patients with renal dysfunction is slightly reduced compared to younger patients with normal renal function. In patients with hepatic diseases, the mean elimination half-life is prolonged by 30 min and the mean plasma clearance is reduced by 1 ml/kg/min. (See also section 4.2).

6. PHARMACEUTICAL PARTICULARS**6.1 List of excipients**

Gluconolactone
Sodium acetate trihydrate
Sodium citrate
Water for injections

6.2 Incompatibilities

Physical incompatibility has been documented for Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion when added to solutions containing the following active substances: amphotericin, amoxicillin, azathioprine, cefazolin, cloxacillin, dexamethasone, diazepam, enoximone, erythromycin, famotidine, furosemide, hydrocortisone sodium succinate, insulin, intralipid, methohexital, methylprednisolone, prednisolone sodium succinate, thiopental, trimethoprim and vancomycin.

This medicinal product must not be mixed with other medicinal products except those mentioned in section 6.6.

6.3 Shelf life

Unopened ampoule: 18 months

After first opening: The product should be used immediately after opening the ampoule.

After dilution:

Chemical and physical in-use stability of a 5.0 mg/ml and 0.1 mg/ml solution (diluted with sodium chloride 9 mg/ml (0.9 %) and glucose 50 mg/ml (5 %) solution for infusion) has been demonstrated for 24 hours at 25 °C exposed to room light in glass, PE and PVC.

From the microbiological point of view, the product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2 to 8 °C, unless dilution has taken place in controlled and validated aseptic conditions.

6.4 Special precautions for storage

Do not store above 25 °C.

For storage conditions after dilution of the medicinal product, see section 6.3.

6.5 Nature and contents of container

Colourless polyethylene LDPE ampoule containing 5 ml. Package sizes:

Packaging of 20 ampoules each containing 5 ml.

6.6 Special precautions for disposal and other handling

Any unused solutions should be discarded.

The solution is to be visually inspected prior to use. Only clear solutions practically free from particles should be used.

Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion has shown to be compatible with: sodium chloride 9 mg/ml (0.9 %) and glucose 50 mg/ml (5 %) solution for infusion, resulting in a dilution of Rocuronium B. Braun of 5 mg/ml and 0.1 mg/ml, respectively.

If Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion is administered via the same infusion line with other medicinal products, it is important that the infusion line is adequately flushed (e.g. with sodium chloride 9 mg/ml (0.9 %) solution for infusion) between administration of Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion and medicinal products for which incompatibility with Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion has been demonstrated or for which compatibility with Rocuronium B. Braun 10 mg/ml Solution for Injection / Infusion has not been established.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

The LDPE ampoules are specially designed to fit Luer lock and Luer fit syringes.

7. PRODUCT REGISTRATION HOLDER

B. Braun Medical Industries 5/B
11900 Bayan Lepas, Penang, Malaysia.

8. DATE OF REVISION OF THE TEXT

10.2024

Product Owner:
B. Braun Melsungen AG,
Carl-Braun-Straße 1,
34212 Melsungen,
Germany

Manufactured by:
B. Braun Melsungen AG,
Mistelweg 2,
12357 Berlin,
Germany

B | BRAUN