

수정사항

1. HOW SUPPLIED 문구 추가

Non-steroidal Anti-inflammatory, Analgesic Agent

ROSIDEN Gel

Piroxicam 5mg

DESCRIPTION : Rosiden Gel is a pale yellow gel in an aluminium tube.

COMPOSITION : Each gram contains 5mg of Piroxicam.

PHARMACODYNAMICS: ROSIDEN is a non-steroidal anti-inflammatory agent which also possesses analgesic and antipyretic properties.

PHARMACOKINETICS: Piroxicam is abosorbed to some degree after topical application.

INDICATIONS : ROSIDEN Gel is indicated in the relief of pain and inflammation of the following disease : arthritis deformans. periomathritis, tendonitis, tendosynovitis, peritendinitis, radiohumeral epicondylitis(tennis elbow etc.), myalgia(myitis, fascitis etc.) and traumatic swelling and alagia.

DOSAGE AND ADMINISTRATION : Apply the ROSIDEN Gel 3 to 4 times daily to the affected areas.

ROUTE OF ADMINISTRATION: Topical

CONTRAINDICATIONS : ROSIDEN Gel is contraindicated in patients with a history of hypersensitivity to Piroxicam.

PREGNANCY AND LACTATION

- a) Pregnancy : Since the safety has not been established in pregnant women, the use of these drugs in pregnancy, nursing mothers, or women of childbearing potential requires that the possible benefits of the drug be weighed against the potential hazards to the mother and embryo or fetus.
- b) Infant : The safety has not been established in infant.

WARNINGS

- a) For external use only. Avoid contact with the eyes and mucosa.
- b) Don't apply this drug to ODT.(Occlusive Dressing Type)
- c) Be careful when apply to the damaged skin, concerning the possibility of temporary irritation.

PRECAUTIONS

- a) ROSIDEN Gel is not for causal treatment but for symptomatic treatment.
- b) For the use of inflammation accompanying infections, carefully use in combination with antibacterial and antifungal agent, concerning the possibility of subclinical infection.
- c) ROSIDEN Gel is not indicated for the treatment of chronic diseases.

INTERACTIONS WITH OTHER MEDICAMENTS: Unknown

SIDE EFFECTS : Eruption may occur occasionally; redness, photosensitivity and pruritus may occur rarely. When these symptoms occur, ROSIDEN Gel should be discontinued immediately. Mild but transient skin discoloration and staining of clothing have been noted when the gel is not rubbed in completely.

SYMPTOMS AND TREATMENT OF OVERDOSE: Unknown

STORAGE: Store in a tight container below 30℃.

HOW SUPPLIED

10 g/Tube, 30 g/Tube, 60 g/ Tube, 100 g/Tube. Not all pack sizes are available in all countries

PRODUCT REGISTRATION HOLDER:THE ZYFAS MEDICAL CO.,
No.7 Jalan Molek 3/10, Taman Molek, 81100 Johor Bahru.

DATE OF REVISION:
June 2023



SHIN POONG PHARM. CO., LTD.
7, Wonsi-ro, Danwon-gu, Ansan-si, Gyeonggi-do, Republic of Korea